Directed Information on Abstract Spaces: Properties and Variational Equalities

Charalambos D. Charalambous and Photios A. Stavrou, *Student Member, IEEE*

arXiv:1302.3971v1 [cs.IT] 16 Feb 2013

Abstract

Directed information or its variants are utilized extensively in the characterization of the capacity of channels with memory and feedback, nonanticipative lossy data compression, and their generalizations to networks.

In this paper, we derive several functional and topological properties of directed information for general abstract alphabets (complete separable metric spaces) using the topology of weak convergence of probability measures. These include convexity of the set of causally conditioned convolutional distributions, convexity and concavity of directed information with respect to sets of such distributions, weak compactness of families of causally conditioned convolutional distributions, their joint distributions and their marginals. Furthermore, we show lower semicontinuity of directed information, and under certain conditions we also establish continuity of directed information. Finally, we derive variational equalities of directed information, which are analogous to those of mutual information (utilized in Blahut-Arimoto algorithm).

In summary, we extend the basic functional and topological properties of mutual information to directed information. Throughout the paper, the importance of the properties of directed information is discussed in the context of extremum problems of directed information.

Index Terms

Directed information, weak convergence, convexity, concavity, lower semicontinuity, continuity, variational equalities.

The authors are with the Department of Electrical and Computer Engineering (ECE), University of Cyprus, 75 Kallipoleos Avenue, P.O. Box 20537, Nicosia, 1678, Cyprus, e-mail: chadcha@ucy.ac.cy., stavrou.fotios@ucy.ac.cy

CONTENTS

I Introduction 3

II Equivalent Nonanticipative Channels on Abstract Spaces 6

III Properties of Directed Information 15

III-A Directed Information Functional of Consistent Conditional Distributions . 18

III-B Convexity and Concavity of Directed Information 22

III-C Weak Convergence and Compactness of Conditional Distributions 25

III-D Lower Semicontinuity of Directed Information 31

III-E Continuity of Directed Information 32

III-F Extension of Directed Information to Arbitrary sequences of RV's 34

IV Variational Equalities of Directed Information 35

V Conclusion 40

VI Appendix A 41

VI-A Backround material 41

VII Appendix B 49

VII-A Additional Discussion 49

VII-A1 Directed Information as a Functional of Sequence of Conditional Distributions 49

VII-A2 Directed Information as the supremum over finite partitions . . 50

VII-B Proof of Theorem III.4 51

VII-C Proof of Theorem III.5 55

VII-D Proof of Theorem III.9 61

References 63

I. INTRODUCTION

Directed information quantifies the directivity of information defined by a causal sequence of feedback and feedforward channel conditional distributions [1], [2]. Specifically, given two sequences of Random Variables (RV's) $X^n \stackrel{\triangle}{=} \{X_0, X_1, \ldots, X_n\} \in \mathcal{X}_{0,n} \stackrel{\triangle}{=} \times_{i=0}^n \mathcal{X}_i$, $Y^n \stackrel{\triangle}{=} \{Y_0, Y_1, \ldots, Y_n\} \in \mathcal{Y}_{0,n} \stackrel{\triangle}{=} \times_{i=0}^n \mathcal{Y}_i$, where $\mathcal{X}_i$ and $\mathcal{Y}_i$ are the input and output alphabets of a channel, respectively, directed information from X^n to Y^n is often defined via conditional mutual information [2], [3]:

$$I(X^n \to Y^n) \stackrel{\triangle}{=} \sum_{i=0}^n I(X^i; Y_i | Y^{i-1}) \tag{I.1}$$

$$= \sum_{i=0}^n \int_{\mathcal{X}_{0,i} \times \mathcal{Y}_{0,i}} \log \left(\frac{P_{Y_i|Y^{i-1},X^i}(dy_i|y^{i-1}, x^i)}{P_{Y_i|Y^{i-1}}(dy_i|y^{i-1})} \right) P_{X^i,Y^i}(dx^i, dy^i) \tag{I.2}$$

$$\equiv \mathbb{I}_{X^n \to Y^n}(P_{X_i|X^{i-1},Y^{i-1}}, P_{Y_i|Y^{i-1},X^i} : i = 0, 1, \ldots, n). \tag{I.3}$$

For each $i = 0, \ldots, n$, $P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot)$ is the conditional distribution of the RV Y_i given the causal information $Y^{i-1} = y^{i-1}$, $X^i = x^i$, $P_{Y_i|Y^{i-1}}(\cdot|\cdot)$ is similarly defined, and $P_{X^i,Y^i}(\cdot,\cdot)$ is the joint distribution of the RV's (X^i, Y^i). Using the decomposition $P_{X^i,Y^i}(dx^i, dy^i) = \otimes_{j=0}^i P_{X_j|X^{j-1},Y^{j-1}} \otimes P_{Y_j|Y^{j-1},X^j}$-a.s., it is clear that directed information $I(X^n \to Y^n)$ is a functional of two collections of causal conditional distributions, the feedforward distribution $\{P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots, n\}$, and the feedback distribution $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots, n\}$, hence the adopted functional notation indicated by (I.3).

In information theory, directed information (I.1) or its variants are used to characterize capacity of channels with memory and feedback [4]–[11], lossy data compression of sequential codes [4], lossy data compression with feedforward information at the decoder [12], lossy data compression of block codes, and capacity of networks, such as, the two-way channel, the multiple access channel [3], [13], etc. Moreover, directed information is also utilized in a variety of problems subject to causality constraints, such as, gambling, portfolio theory, data compression and hypothesis testing [14], and in biology as an alternative to Granger's measure of causality [15]–[17]. Some of the above references derive coding theorems under any of the assumptions: (a) stationary ergodic processes $\{(X_i, Y_i) : i = 0, 1, \ldots\}$, (b) Dobrushin's stability of the information density $\frac{1}{n+1} \log \otimes_{i=0}^n \frac{P_{Y_i|Y^{i-1},X^i}(dy_i|y^{i-1},x^i)}{P_{Y_i|Y^{i-1}}(dy_i|y^{i-1})}$, or (c) its generalization using the information spectrum methods [18]. Hence, relations between $I(X^n \to Y^n)$ or its information spectrum to the optimal

rates theoretically achievable are established in an anthology of problems of information theory.

Directed information as initially introduced by Marko [1] is obtained via a decomposition of Shannon's self-mutual information into two directional parts, and then taking their expectations. Although directed information is obtained from mutual information, its functional and topological properties are not well understood [3], compared to those of mutual information. Specific functional properties of mutual information expressed as a functional $I(X^n;Y^n) \equiv \mathbb{I}_{X^n;Y^n}(P_{X^n}, P_{Y^n|X^n})$, of the two distributions $\{P_{X^n}, P_{Y^n|X^n}\}$, such as, convexity, concavity, and topological properties such as lower semicontinuity (with respect to the topology of weak convergence of probability measures), at first glance, do not translate into analogous properties for directed information $I(X^n \to Y^n) \equiv \mathbb{I}_{X^n \to Y^n}(P_{X_i|X^{i-1},Y^{i-1}}, P_{Y_i|Y^{i-1},X^i} : i = 0, 1, \ldots, n)$, as a functional of the two distributions $\{P_{X_i|X^{i-1},Y^{i-1}}, P_{Y_i|Y^{i-1},X^i} : i = 0, 1, \ldots, n\}$. Similarly, it is not obvious whether the well-known variational equalities of mutual information utilized in the Blahut-Arimoto algorithm [19], can be extended to directed information. These properties together with compactness of subsets of the sets of the conditional distributions $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots, n\}$ and $\{P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots, n\}$, are fundamental in addressing extremum problems of directed information related to channel capacity, sequential and nonanticipative rate distortion, and their generalizations to networks, for abstract (e.g., continuous) alphabets. In fact, as shown in [20] even the analysis of single letter mutual information defined on abstract alphabets becomes very technical, and hence the same is expected for directed information.

The main objective of this paper is to determine whether the functional and topological properties of mutual information can be extended to corresponding properties of directed information defined on abstract Polish spaces (complete separable metric spaces), and to provide appropriate conditions for these extensions to hold. The main results are summarized below.

R1) Introduce an equivalent directed information definition expressed via information divergence $\mathbb{D}(\cdot||\cdot)$, as a functional of two consistent families of conditional distributions $\mathbf{P}(\cdot|\mathbf{y})$ on $\mathcal{X}^{\mathbb{N}} \stackrel{\triangle}{=} \times_{i=0}^{\infty}\mathcal{X}_i$ for $\mathbf{y} = (y_0, y_1, \ldots) \in \mathcal{Y}^{\mathbb{N}} \stackrel{\triangle}{=} \times_{i=0}^{\infty}\mathcal{Y}_i$, and $\mathbf{Q}(\cdot|\mathbf{x})$ on $\mathcal{Y}^{\mathbb{N}}$ for $\mathbf{x} \in \mathcal{X}^{\mathbb{N}}$, which uniquely define $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots\}$ and $\{P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots\}$, respectively, and vice-versa, and their $(n+1)$-fold convolutional measures $\overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \triangleq$

$\otimes_{i=0}^{n} P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot)$, $\overrightarrow{Q}_{0,n}(dy^n|x^n) \triangleq \otimes_{i=0}^{n} P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot)$;

R2) Show convexity of the consistent families of the conditional distributions $\mathbf{P}(\cdot|\mathbf{y})$, $\mathbf{Q}(\cdot|\mathbf{x})$ as subsets of the set of regular conditional distributions;

R3) Show convexity and concavity of directed information as a functional with respect to the consistent families of conditional distributions $\mathbf{Q}(\cdot|\mathbf{x})$ and $\mathbf{P}(\cdot|\mathbf{y})$, respectively;

R4) Show weak compactness of the consistent families of conditional distributions $\mathbf{P}(\cdot|\mathbf{x})$ and $\mathbf{Q}(\cdot|\mathbf{y})$, and of their marginals and joint distribution;

R5) Show lower semicontinuity of directed information as a functional of the consistent families of the conditional distributions $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$, and continuity of directed information as a functional of the family $\mathbf{P}(\cdot|\mathbf{y})$;

R6) Express directed information in terms of variational equalities involving minimization and maximization operations over consistent families of conditional distributions;

R7) Illustrate that the functional and topological properties obtained in R1)–R6) extend naturally to three sequences of RV's $X^n \in \mathcal{X}_{0,n}$, $Y^n \in \mathcal{Y}_{0,n}$, $Z^n \in \mathcal{Z}_{0,n}$, or more, which cover directed information measures for networks, and problems with side information.

The above functional and topological properties are shown by invoking the topology of weak convergence of probability measures on Polish spaces and Prohorov's theorems. Some of the results described above are obtained by utilizing some analogies between communication channels with memory and feedback, and stochastic control in which the control process and the controlled process are sequences of conditional distributions, which are analogous to $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots\}$ and $\{P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots\}$, respectively, [21], [22].

From the practical point of view, there are many potential applications of the functional and topological properties of directed information derived in this paper. Below we list some of them. The concavity and convexity properties are important in deriving tight bounds for converse coding theorems, and in identifying properties of extremum problems involving feedback capacity [3], [23] and sequential and nonanticipative lossy data compression for point-to-point [24], and network communication [25], [26]. The semicontinuity and continuity of directed information, and the compactness of the consistent families of distributions $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$ are crusial when addressing questions of existence of extremum solutions to problems involving feedback capacity, sequential and nonanticipative lossy data compression, computations of extremum

solutions and their properties, and derivations of coding theorems on abstract alphabets. The point to be made is that R1)–R5) are fundamental properties in any extremum problem involving directed information. The variational equalities are important in generalizing the Blahut-Arimoto computation schemes to information measures involving directed information.

For the sake of clarity, we discuss one application of the results. Consider the extremum problem of channel capacity with memory and feedback. Under the assumption of stationary ergodic processes $\{(X_i, Y_i) : i = 0, 1, \ldots\}$ or Dobrushin's information stability, the operational definition of capacity is given by the following extremum problem [8].

$$C^f(P) \stackrel{\triangle}{=} \liminf_{n\to\infty} \sup_{\left\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot): i=0,1,\ldots,n\right\}\in\mathcal{P}_{0,n}(P)} \frac{1}{n+1} I(X^n \to Y^n), \tag{I.4}$$


where $\mathcal{P}_{0,n}(P)$ denotes the power constraint set. The task of showing existence of a measure $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots, n\} \in \mathcal{P}_{0,n}(P)$ which achieves the supremum in (I.4) is not easy. The main difficulty arises from the fact that $I(X^n \to Y^n)$ is a functional of the two sequences of distributions $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot), P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots, n\}$, unlike mutual information $I(X^n; Y^n) \equiv \mathbb{I}_{X^n;Y^n}(P_{X^n}, P_{Y^n|X^n})$, which inherits most of its properties from those of relative entropy between two distributions. Even for problems involving mutual information, Csiszár [20], [27] introduced elaborate arguments to address such extremum problems on abstract alphabets. However, by utilizing R1)–R5) it is possible to show existence of such conditional distribution not only for (I.4), but also for the sequential and nonanticipative rate distortion functions, obtain generalizations involving three or more random processes, and identify the properties of the extremum solutions.

The rest of the paper is structured as follows. Section II introduces two equivalent definitions of nonanticipative channels on abstract spaces $\big($R1)$\big)$. Section III derives the functional and topological properties of directed information $\big($R2)–R5)$\big)$. Section IV derives variational equalities of directed information $\big($R6)$\big)$. Finally, Section V gives concluding remarks and identifies additional problems of future interest. Appendix VII gives a brief discussion of the backround material utilized to derive the main results and includes some of the proofs.

II. Equivalent Nonanticipative Channels on Abstract Spaces

In this section, our aim is to establish two equivalent definitions of the sequence of conditional distributions or basic processes, which define any probabilistic channel with nonanticipative

(causal) feedback, that relate causally the input-output behavior of the channel. This formulation is necessary to establish the results stated under R1)–R7). The first definition of conditional distributions is the usual one found in many papers, e.g., [3], [4], [7]–[10]; it is described via a family of functions which are regular conditional distributions so that their convolution product is a family of regular conditional distributions on product alphabets (see Fig. II.1, (a)). The second definition is described via a family of regular conditional distributions defined on product alphabets, which satisfy a certain consistency condition (see Fig. II.1, (b)). The



(a) Sequence of feedback and feedforward channels $\{P_{X_i|X^{i-1},Y^{i-1}}, P_{Y_i|Y^{i-1},X^i} : i = 0, 1, \ldots, n\}$.

(b) Consistent families of feedback and feedforward channels $\{\overleftarrow{P}_{X^n|Y^{n-1}}, \overrightarrow{Q}_{Y^n|X^n} : n \in \mathbb{N}\}$.

Fig. II.1. Equivalent Representations of Feedback/Feedforward Channels.

second definition is often utilized in the stochastic control literature, when both the control and controlled processes are conditional probability distributions [21], [22]. Indeed, the analogy is that $\{X_i : \ i = 0, 1, \ldots\}$ is the controlled process and $\{Y_i : \ i = 0, 1, \ldots\}$ is the control process. The second definition is appealing from the point of view of expressing the information density $i(X^n \to Y^n) \stackrel{\triangle}{=} \log \otimes_{i=0}^{n} \frac{P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot)}{P_{Y_i|Y^{i-1}}(\cdot|\cdot)}$ associated with directed information $I(X^n \to Y^n)$, in terms of two consistent families of conditional distributions, namely, $\mathbf{Q}(\cdot|\mathbf{x})$ on $\mathcal{Y}^{\mathbb{N}}$ given $\mathbf{x} = (x_0, x_1, \ldots) \in \mathcal{X}^{\mathbb{N}}$, and $\mathbf{P}(\cdot|\mathbf{y})$ on $\mathcal{X}^{\mathbb{N}}$ given $\mathbf{y} = (y_0, y_1, \ldots) \in \mathcal{Y}^{\mathbb{N}}$, which uniquely define $\{P_{Y_i|Y^{i-1},X^i}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots\}$ and $\{P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot) : i = 0, 1, \ldots\}$, respectively, such that $i(X^n \to Y^n) = \log \frac{\mathbf{Q}(dy^n|x^n)}{\nu^{\mathbf{P}\otimes\mathbf{Q}}(dy^n)} - a.s.$, where $\nu^{\mathbf{P}\otimes\mathbf{Q}}(\cdot)$ is the marginal distribution[1] on $\times_{i=0}^{n}\mathcal{Y}_i$ obtained from $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$. Once the conditions on the abstract spaces $\{(\mathcal{Y}_i, \mathcal{X}_i) : \ i = 0, 1, \ldots\}$ are identified, and the consistency conditions are introduced, then it can be shown that $i(X^n \to Y^n)$ has another version given by $i(X^n \to Y^n) = \log \frac{\mathbf{P}(\cdot|\cdot)\otimes\mathbf{Q}(\cdot|\cdot)}{\mathbf{P}(\cdot|\cdot)\otimes\nu^{\mathbf{P}\otimes\mathbf{Q}}(\cdot)}(x^n, y^n) - a.s.$, where $\otimes$ denotes convolution of measures. Consequently, directed information can be expressed

[1]In the rest of the paper we write ν instead of $\nu^{\mathbf{P}\otimes\mathbf{Q}}$ omitting its explicit dependence on $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$.

in terms of Kullback-Leibler distance $\mathbb{D}\big(\frac{\mathbf{P}(\cdot|\cdot)\otimes\mathbf{Q}(\cdot|\cdot)}{\mathbf{P}(\cdot|\cdot)\otimes\nu^{\mathbf{P}\otimes\mathbf{Q}}(\cdot)}\big)$. Results R1)–R6) are obtained by utilizing the second equivalent definition of conditional distributions.

Notations and Preliminaries.

Denote the set of non-negative integers by $\mathbb{N} \stackrel{\triangle}{=} \{0,1,2,\ldots\}$, and its restriction to a finite set by $\mathbb{N}^n \stackrel{\triangle}{=} \{0,1,2,\ldots,n\}$. Introduce two sequence of spaces $\{(\mathcal{X}_n, \mathcal{B}(\mathcal{X}_n)) : n \in \mathbb{N}\}$ and $\{(\mathcal{Y}_n, \mathcal{B}(\mathcal{Y}_n)) : n \in \mathbb{N}\}$, called basic measurable spaces, where $\mathcal{X}_n, \mathcal{Y}_n, n \in \mathbb{N}$ are topological spaces, and $\mathcal{B}(\mathcal{X}_n)$ and $\mathcal{B}(\mathcal{Y}_n)$ are Borel $\sigma-$algebras of subsets of $\mathcal{X}_n$ and $\mathcal{Y}_n$, respectively.

For each $n \in \mathbb{N}$ define the product spaces

$$(\mathcal{X}_{0,n}, \mathcal{B}(\mathcal{X}_{0,n})) \stackrel{\triangle}{=} (\times_{i=0}^{n}\mathcal{X}_i, \odot_{i=0}^{n}\mathcal{B}(\mathcal{X}_i)), (\mathcal{Y}_{0,n}, \mathcal{B}(\mathcal{Y}_{0,n})) \stackrel{\triangle}{=} (\times_{i=0}^{n}\mathcal{Y}_i, \odot_{i=0}^{n}\mathcal{B}(\mathcal{Y}_i)).$$

The basic measurable spaces are connected to a random experiment consisting of a countable chain of trials with a time ordering as follows. For each $n \in \mathbb{N}$, let $\mathcal{X}_n$ and $\mathcal{Y}_n$ be the spaces of all possible outcomes at time $n \in \mathbb{N}$. Having performed the trials up to and including the nth time, the time ordering of outcomes is $x_0, y_0, x_1, y_1, \ldots, x_n, y_n$ in $\mathcal{X}_0, \mathcal{Y}_0, \mathcal{X}_1, \mathcal{Y}_1, \ldots, \mathcal{X}_n, \mathcal{Y}_n$, respectively. The probability distributions governing the next trial at time $(n+1)$ are $p_{n+1}(A_{n+1}; x_0, \ldots, x_n, y_0, \ldots, y_n)$ and $q_{n+1}(B_{n+1}; y_0, \ldots, y_n, x_0, \ldots, x_{n+1})$, $A_{n+1} \in \mathcal{B}(\mathcal{X}_{n+1})$, $B_{n+1} \in \mathcal{B}(\mathcal{Y}_{n+1})$. Hence, each possible outcome of the experiment is a sequence $\omega = (x_0, y_0, x_1, y_1, \ldots)$ with $x_n \in \mathcal{X}_n, y_n \in \mathcal{Y}_n$ for each $n \in \mathbb{N}$.

Consequently, define the sample space Ω and the algebra $\mathcal{F}$ of all experiments by

$$(\Omega, \mathcal{F}) \stackrel{\triangle}{=} \Big(\times_{n\in\mathbb{N}} (\mathcal{X}_n \times \mathcal{Y}_n), \odot_{n\in\mathbb{N}}\mathcal{B}(\mathcal{X}_n) \odot \mathcal{B}(\mathcal{Y}_n)\Big).$$

Associated with the basic measurable spaces there are two basic sequences of Random Variables (RV's) $\{X_n : n \in \mathbb{N}\}$ and $\{Y_n : n \in \mathbb{N}\}$, such that for each $n \in \mathbb{N}$, they take values $X_n \in \mathcal{X}_n$ and $Y_n \in \mathcal{Y}_n$. These are introduced as follows.

Let $X_0, Y_0, X_1, Y_1, \ldots$ be the coordinate RV's. For each $n \in \mathbb{N}$

$$X_n(\omega) = x_n, \;\; Y_n(\omega) = y_n \;\; \text{ if } \;\; \omega = (x_0, y_0, x_1, y_1, \ldots).$$

Clearly, $X_n : (\Omega, \mathcal{F}) \mapsto (\mathcal{X}_n, \mathcal{B}(\mathcal{X}_n))$, $Y_n : (\Omega, \mathcal{F}) \mapsto (\mathcal{Y}_n, \mathcal{B}(\mathcal{Y}_n))$, and for each outcome $\omega \in \Omega$ of the experiment, $X_n(\omega)$, $Y_n(\omega)$ are the results of the nth time. Similarly, $X^n \stackrel{\triangle}{=} \{X_0, \ldots, X_n\}$ and $Y^n \stackrel{\triangle}{=} \{Y_0, \ldots, Y_n\}$ denote the result of the trials up to and including the nth time; they are RV taking values in $(\mathcal{X}_{0,n}, \mathcal{B}(\mathcal{X}_{0,n}))$ and $(\mathcal{Y}_{0,n}, \mathcal{B}(\mathcal{Y}_{0,n}))$, respectively. The objective is to

construct a measure $\mathbb{P}$ on $(\Omega, \mathcal{F})$ consistent with the data (e.g., measurable spaces and conditional distributions).

For every $n \in \mathbb{N}$, define the σ-algebras generated by $\{X_0, X_1, \ldots, X_n\}$ and $\{Y_0, Y_1, \ldots, Y_n\}$ by

$$\mathcal{F}(X^n) \stackrel{\triangle}{=} \sigma\{X_0, X_1, \ldots, X_n\},\ \mathcal{F}(Y^n) \stackrel{\triangle}{=} \sigma\{Y_0, Y_1, \ldots, Y_n\}.$$

Then every event $H \in \mathcal{F}(X^n)$ has the form

$$H = \Big\{(X_0, X_1, \ldots, X_n) \in A\Big\} = A \times \mathcal{X}_{n+1} \times \ldots \times \ldots,\ A \in \mathcal{B}(\mathcal{X}_{0,n})$$

and H is called a cylinder set with base $A \in \mathcal{B}(\mathcal{X}_{0,n})$. Similarly, for an event $J \in \mathcal{F}(Y^n)$

$$J = \Big\{(Y_0, Y_1, \ldots, Y_n) \in B\Big\} = B \times \mathcal{Y}_{n+1} \times \ldots \times \ldots,\ B \in \mathcal{B}(\mathcal{Y}_{0,n})$$

and J is a cylinder set with base $B \in \mathcal{B}(\mathcal{Y}_{0,n})$.

Points in the Cartesian countable product spaces $\mathcal{X}^{\mathbb{N}} \stackrel{\triangle}{=} \times_{n\in\mathbb{N}}\mathcal{X}_n$, $\mathcal{Y}^{\mathbb{N}} \stackrel{\triangle}{=} \times_{n\in\mathbb{N}}\mathcal{Y}_n$ are denoted by $\mathbf{x} \stackrel{\triangle}{=} \{x_0, x_1, \ldots\} \in \mathcal{X}^{\mathbb{N}}$, $\mathbf{y} \stackrel{\triangle}{=} \{y_0, y_1, \ldots\} \in \mathcal{Y}^{\mathbb{N}}$, respectively. Similarly, for $n \in \mathbb{N}$, points in $\mathcal{X}_{0,n} \stackrel{\triangle}{=} \times_{i=0}^{n}\mathcal{X}_i$, $\mathcal{Y}_{0,n} \stackrel{\triangle}{=} \times_{i=0}^{n}\mathcal{Y}_i$ are denoted by $x^n \stackrel{\triangle}{=} \{x_0, x_1, \ldots, x_n\} \in \mathcal{X}_{0,n}$, $y^n \stackrel{\triangle}{=} \{y_0, y_1, \ldots, y_n\} \in \mathcal{Y}_{0,n}$, respectively.

Let $\mathcal{B}(\mathcal{X}^{\mathbb{N}})$ and $\mathcal{B}(\mathcal{Y}^{\mathbb{N}})$ denote the $\sigma-$algebras in $\mathcal{X}^{\mathbb{N}}$ and $\mathcal{Y}^{\mathbb{N}}$, respectively, generated by cylinder sets (e.g., $\mathcal{B}(\mathcal{X}^{\mathbb{N}})$ is the smallest Borel $\sigma-$algebra containing all cylinder sets $\{\mathbf{x} = (x_0, x_1, \ldots) \in \mathcal{X}^{\mathbb{N}} : x_0 \in A_0, x_1 \in A_1, \ldots, x_n \in A_n\}, A_i \in \mathcal{B}(\mathcal{X}_i), 0 \le i \le n, n \ge 0$). The Borel σ-algebra $\mathcal{B}(\mathcal{X}^{\mathbb{N}})$ is denoted by $\odot_{i\in\mathbb{N}}\mathcal{B}(\mathcal{X}_i)$. Hence, $\mathcal{B}(\mathcal{X}_{0,n})$ and $\mathcal{B}(\mathcal{Y}_{0,n})$ denote the $\sigma-$algebras of cylinder sets in $\mathcal{X}^{\mathbb{N}}$ and $\mathcal{Y}^{\mathbb{N}}$, respectively, with bases over $A_i \in \mathcal{B}(\mathcal{X}_i),\ i = 0, 1, \ldots, n$, and $B_i \in \mathcal{B}(\mathcal{Y}_i),\ i = 0, 1, \ldots, n$, respectively.

Backward or Feedback Channel.

Suppose for each $n \in \mathbb{N}$, the conditional distribution of the RV $X_n \in \mathcal{X}_n$ is determined provided the values of the basic processes $X^{n-1} = x^{n-1} \in \mathcal{X}_{0,n-1}$ and $Y^{n-1} = y^{n-1} \in \mathcal{Y}_{0,n-1}$ are known, and let $\{p_n(dx_n; x^{n-1}, y^{n-1}) : n \in \mathbb{N}\}$ denote this family of distributions. Here, it is assumed that $\sigma\{X^{-1}, Y^{-1}\} = \{\emptyset, \Omega\}$, hence $p_0(dx_0; x^{-1}, y^{-1}) \stackrel{\triangle}{=} p_0(x_0)$. The functions $p_n(\cdot; \cdot, \cdot)$ are candidates of distributions of the sequence of RV's $\{X_n : n \in \mathbb{N}\}$ on $\{(\mathcal{X}_n, \mathcal{B}(\mathcal{X}_n)) : n \in \mathbb{N}\}$ if and only if the following conditions hold.

i) For every $n \in \mathbb{N}$, $p_n(\cdot; x^{n-1}, y^{n-1})$ is a probability measure on $\mathcal{B}(\mathcal{X}_n)$;

ii) For every $n \in \mathbb{N}$, $p_n(A_n; x^{n-1}, y^{n-1})$ is a $\odot_{i=0}^{n-1}\big(\mathcal{B}(\mathcal{X}_i) \odot \mathcal{B}(\mathcal{Y}_i)\big)$-measurable function of

$x^{n-1} \in \mathcal{X}_{0,n-1}$, $y^{n-1} \in \mathcal{Y}_{0,n-1}$, for every $A_n \in \mathcal{B}(\mathcal{X}_n)$.

Given the collection of functions $\{p_n(dx_n; x^{n-1}, y^{n-1}) : n \in \mathbb{N}\}$ satisfying conditions **i), ii)**, one can construct a family of distributions on the product space $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})) \stackrel{\triangle}{=} \left(\times_{i\in\mathbb{N}}\mathcal{X}_i, \odot_{i\in\mathbb{N}}\mathcal{B}(\mathcal{X}_i)\right)$ as follows.

Let $C \in \mathcal{B}(\mathcal{X}_{0,n})$ be a cylinder set of the form

$$C \stackrel{\triangle}{=} \Big\{\mathbf{x} \in \mathcal{X}^{\mathbb{N}} : x_0 \in C_0, x_1 \in C_1, \ldots, x_n \in C_n\Big\},\ C_i \in \mathcal{B}(\mathcal{X}_i),\ 0 \le i \le n.$$

Define a family of measures $\mathbf{P}(\cdot|\mathbf{y})$ on $\mathcal{B}(\mathcal{X}^{\mathbb{N}})$ by

$$\mathbf{P}(C|\mathbf{y}) \stackrel{\triangle}{=} \int_{C_0} p_0(dx_0) \int_{C_1} p_1(dx_1; x^0, y^0) \ldots \int_{C_n} p_n(dx_n; x^{n-1}, y^{n-1}) \tag{II.1}$$

$$\equiv \overleftarrow{P}_{0,n}(C_{0,n}|y^{n-1}),\ C_{0,n} = \times_{i=0}^{n} C_i. \tag{II.2}$$

The notation $\overleftarrow{P}_{0,n}(C_{0,n}|y^{n-1})$ is used to denote the causal conditioning dependence of the measure $\mathbf{P}(\cdot|\mathbf{y})$ defined on cylinder sets $C \in \mathcal{B}(\mathcal{X}_{0,n})$, for any $n \in \mathbb{N}$. The right hand side of (II.1) uniquely defines a measure on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$. Moreover, for each $n \in \mathbb{N}$ the family of measures $\mathbf{P}(\cdot|\mathbf{y})$ satisfies the following property (inherited from condition **ii)**): for $E \in \mathcal{B}(\mathcal{X}^{\mathbb{N}})$, $\mathbf{P}(E|\mathbf{y})$ is $\mathcal{B}(\mathcal{Y}^{\mathbb{N}})-$measurable, and for $E \in \mathcal{B}(\mathcal{X}_{0,n})$, $\mathbf{P}(E|\mathbf{y})$ is $\mathcal{B}(\mathcal{Y}_{0,n-1})-$measurable.

Thus, if conditions **i)** and **ii)** hold then for each $\mathbf{y} \in \mathcal{Y}^{\mathbb{N}}$, the right hand side of (II.1) defines a consistent family of finite-dimensional distribution, and hence there exists a unique measure on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$, for which $p_n(dx_n; x^{n-1}, y^{n-1})$ is obtained. This leads to the first definition of a feedback channel, as a family of functions $p_n(dx_n; x^{n-1}, y^{n-1})$ satisfying conditions **i)** and **ii)**. This definition is used extensively by many authors [3], [4], [7]–[10].

An alternative, equivalent definition of a feedback channel is established as follows. Consider a family of measures $\mathbf{P}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ satisfying the following consistency condition.

C1: If $E \in \mathcal{B}(\mathcal{X}_{0,n})$ then $\mathbf{P}(E_{0,n}|\mathbf{y})$ is $\mathcal{B}(\mathcal{Y}_{0,n-1})-$measurable function of $\mathbf{y} \in \mathcal{Y}^{\mathbb{N}}$.

Clearly, if conditions **i)** and **ii)** are satisfied then **C1** holds for the family of measures $\mathbf{P}(\cdot|\mathbf{y})$ defined via the right hand side of (II.1). The question we address next is whether for any family of measures $\mathbf{P}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ satisfying consistency condition **C1**, one can construct a collection of functions $\{p_n(dx_n; x^{n-1}, y^{n-1}) : n \in \mathbb{N}\}$ satisfying conditions **i)** and **ii)**, which are connected to $\mathbf{P}(\cdot|\cdot)$ via relation (II.1). To illustrate this point, let $A^{(n)} = \{\mathbf{x} \in \mathcal{X}^{\mathbb{N}} : x_n \in A\}$,

$A \in \mathcal{B}(\mathcal{X}_n)$, and let $\mathbf{P}(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1}))$ denote the conditional probability of $A^{(n)}$ with respect to $\mathcal{B}(\mathcal{X}_{0,n-1})$ calculated on the probability space $\big(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}), \mathbf{P}(\cdot|\mathbf{y})\big)$. Then

$$\mathbf{P}(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1})) = p_n(A; x^{n-1}, y^{n-1}), \quad A^{(n)} \in \mathcal{B}(\mathcal{X}_{0,n}), \tag{II.3}$$

for $\mathbf{P}(\cdot|\mathbf{y})-$almost all $\mathbf{x} \in \mathcal{X}^{\mathbb{N}}$. Clearly, the function on the right hand side of (II.3), $p_n(A; x^{n-1}, y^{n-1})$ is $\mathcal{B}(\mathcal{X}_{0,n-1})-$measurable for a fixed $A \in \mathcal{B}(\mathcal{X}_n)$ and $y^{n-1} \in \mathcal{Y}_{0,n-1}$, but it cannot be claimed that $p_n(\cdot; x^{n-1}, y^{n-1})$ is a probability measure on $\mathcal{X}_n$. However, under the general assumption that $\{(\mathcal{X}_n, \mathcal{B}(\mathcal{X}_n)) : n \in \mathbb{N}\}$ are complete separable metric spaces (Polish spaces), with $\mathcal{B}(\mathcal{X}_n)$ the $\sigma-$algebra of Borel sets it is shown in [21] (Appendix VI-A, Theorem VI.16), that the right hand side of (II.3) represents a version of conditional probability $(a.s.)$ such that condition **i)** holds as well. Therefore, to establish the second equivalent definition, introduce the following condition on the alphabet spaces.

iii) $\{\mathcal{X}_n : n \in \mathbb{N}\}$ are complete separable metric spaces and $\{\mathcal{B}(\mathcal{X}_n) : n \in \mathbb{N}\}$ are the $\sigma-$algebras of Borel sets.

By Theorem VI.16, Appendix VI-A, if condition **iii)** holds, then for any family of measures $\mathbf{P}(\cdot|\mathbf{y})$ satisfying **C1** one can construct a collection of versions of conditional distributions $\{p_n(dx_n; x^{n-1}, y^{n-1}) : n \in \mathbb{N}\}$ satisfying conditions **i)** and **ii)** which are connected with $\mathbf{P}(\cdot|\mathbf{y})$ via relation (II.1), and hence the following conclusion. When $\{\mathcal{X}_n : n \in \mathbb{N}\}$ are Polish Spaces with $\{\mathcal{B}(\mathcal{X}_n) : n \in \mathbb{N}\}$ the $\sigma-$algebra of Borel sets, there are two equivalent definitions of a feedback channel. The first definition is the usual one given by a family of functions $p_n(\cdot; x^{n-1}, y^{n-1})$ satisfying conditions **i)** and **ii)**. The second definition is given by a family of measures $\mathbf{P}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ depending parametrically on $\mathbf{y} \in \mathcal{Y}^{\mathbb{N}}$ and satisfying the consistency condition **C1**. Although, the family of measures $\mathbf{P}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ are finite additive probability measures, by Kolmogorov's extension theorem [28], the completeness of $\{\mathcal{X}_n : n \in \mathbb{N}\}$ guarantees the existence of countable additive probability measures $\mathbf{P}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$, whose marginal on each $\mathcal{X}_{0,n}$ is $\overleftarrow{P}_{0,n}(\cdot|y^{n-1})$.

The second equivalent definition of a feedback channel, together with an analogous equivalent similar definition for the forward channel will be used throughout the paper to derive the functional and topological properties of directed information.

Feedforward Channel.

The previous methodology is repeated to obtain two equivalent definitions for the forward channel as well. Suppose for each $n \in \mathbb{N}$, the conditional distribution of the RV $Y_n \in \mathcal{Y}_n$ is determined provided the values of the basic processes $Y^{n-1} \in \mathcal{Y}_{0,n-1}$ and $X^n = x^n \in \mathcal{X}_{0,n}$ are known, and let $\{q_n(dy_n; y^{n-1}, x^n) : n \in \mathbb{N}\}$ denotes this family of distributions. Similarly define $q_0(dy_0; y^{-1}, x^0) \stackrel{\triangle}{=} q_0(dy_0; x_0)$. The collection of functions $\{q_n(dy_n; y^{n-1}, x^n) : n \in \mathbb{N}\}$ satisfy the following conditions.

iv) For every $n \in \mathbb{N}$, $q_n(\cdot; y^{n-1}, x^n)$ is a probability measure $\mathcal{B}(\mathcal{Y}_n)$;
v) For every $n \in \mathbb{N}$, $q_n(B_n; y^{n-1}, x^n)$ is a $\odot_{i=0}^{n-1}\big(\mathcal{B}(\mathcal{Y}_i) \odot \mathcal{B}(\mathcal{X}_i)\big) \odot \mathcal{B}(\mathcal{X}_n)$-measurable function of $x^n \in \mathcal{X}_{0,n}$, $y^{n-1} \in \mathcal{Y}_{0,n-1}$, for every $B_n \in \mathcal{B}(\mathcal{Y}_n)$.

Similarly as before, using the collection of functions $\{q_n(\cdot;\cdot,\cdot) : n \in \mathbb{N}\}$ one can construct a family of measures $\mathbf{Q}(\cdot|\mathbf{x})$ on $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ which depend parametrically on $\mathbf{x} \in \mathcal{X}^{\mathbb{N}}$, as follows. Consider a cylinder set $D \in \mathcal{B}(\mathcal{Y}_{0,n})$ of the form

$$D \stackrel{\triangle}{=} \Big\{\mathbf{y} \in \mathcal{Y}^{\mathbb{N}} : y_0 \in D_0, y_1 \in D_1, \ldots, y_n \in D_n\Big\},\ D_i \in \mathcal{B}(\mathcal{Y}_i),\ 0 \le i \le n.$$

Define a family of measures on $\mathcal{B}(\mathcal{Y}^{\mathbb{N}})$ by

$$\mathbf{Q}(D|\mathbf{x}) \stackrel{\triangle}{=} \int_{D_0} q_0(dy_0; x_0) \int_{D_1} q_1(dy_1; y_0, x^1) \ldots \int_{D_n} q_n(dy_n; y^{n-1}, x^n) \tag{II.4}$$

$$\equiv \overrightarrow{Q}_{0,n}(D_{0,n}|x^n),\ D_{0,n} = \times_{i=0}^{n} D_i. \tag{II.5}$$

Since, for each $\mathbf{x} \in \mathcal{X}^{\mathbb{N}}$ the right hand side of (II.4) defines a consistent family of finite dimensional distribution, then there exist a unique measure on $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ for which the family of distributions $\{q_n(dy_n; y^{n-1}, x^n) : n \in \mathbb{N}\}$ is satisfied. Moreover, the family of measures $\mathbf{Q}(D|\mathbf{x})$ satisfies the following consistency condition.

C2: If $F \in \mathcal{B}(\mathcal{Y}_{0,n})$, then $\mathbf{Q}(F|\mathbf{x})$ is a $\mathcal{B}(\mathcal{X}_{0,n})-$measurable function of $x \in \mathcal{X}^{\mathbb{N}}$.

By Theorem VI.16, Appendix VI-A, to obtain another equivalent definition for the forward channel introduce the following condition on the output alphabet.

vi) $\{\mathcal{Y}_n : n \in \mathbb{N}\}$ are Polish Spaces and $\{\mathcal{B}(\mathcal{Y}_n) : n \in \mathbb{N}\}$ are the $\sigma-$algebra of Borel sets.

Then for any family of measures $\mathbf{Q}(\cdot|\mathbf{x})$ on $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ satisfying consistency condition **C2** $\big($under condition **vi**$\big)$, one can construct a collection of functions $\{q_n(\cdot;\cdot,\cdot) : n \in \mathbb{N}\}$ satisfying conditions **iv)** and **v)**, which are connected with $\mathbf{Q}(\cdot|\mathbf{x})$ via relation (II.4). Therefore, we arrive

at two equivalent definitions for the forward channel as well.

We conclude this section by constructing the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, as stated earlier, and the sequence of RV's $\{(X_i, Y_i) : \ i = 0, 1, \ldots, n\}$ defined on it. Given the basic measures $\mathbf{P}(\cdot|\mathbf{y})$ on $\mathcal{X}^{\mathbb{N}}$ satisfying consistency condition **C1** and $\mathbf{Q}(\cdot|\mathbf{x})$ on $\mathcal{Y}^{\mathbb{N}}$ satisfying consistency condition **C2**, one can construct a sequence of RV's $\{X_n, Y_n : n \in \mathbb{N}\}$ or conditional distributions as follows.

Let $A^{(n)} = \{\mathbf{x} : x_n \in A\}$, $A \in \mathcal{B}(\mathcal{X}_n)$ and $B^{(n)} = \{\mathbf{y} : y_n \in B\}$, $B \in \mathcal{B}(\mathcal{Y}_n)$. In addition, let $\mathbf{P}(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1}))$ denote the conditional probability of $A^{(n)}$ with respect to $\mathcal{B}(\mathcal{X}_{0,n-1})$ calculated on the probability space $\big(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}), \mathbf{P}(\cdot|\mathbf{y})\big)$, and $\mathbf{Q}(B^{(n)}|\mathbf{x}|\mathcal{B}(\mathcal{Y}_{0,n-1}))$ denote the conditional probability of $B^{(n)}$ with respect to $\mathcal{B}(\mathcal{Y}_{0,n-1})$ calculated on the probability space $\big(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \mathbf{Q}(\cdot|\mathbf{x})\big)$.

Then by conditioning it follows that

$$\begin{aligned}\mathbb{P}\big\{X_n \in A_n | X^{n-1} = x^{n-1}, Y^{n-1} = y^{n-1}\big\} &= \mathbf{P}\big(\{\mathbf{x} : x_n \in A_n\}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1})\big), \ A_n \in \mathcal{B}(\mathcal{X}_n) \\ &= p_n(A_n; x^{n-1}, y^{n-1}) \qquad \text{(II.6)} \\ \mathbb{P}\big\{Y_n \in B_n | Y^{n-1} = y^{n-1}, X^n = x^n\big\} &= \mathbf{Q}\big(\{\mathbf{y} : y_n \in B_n\}|\mathbf{x}|\mathcal{B}(\mathcal{Y}_{0,n-1})\big), \ B_n \in \mathcal{B}(\mathcal{Y}_n) \\ &= q_n(B_n; y^{n-1}, x^n) \qquad \text{(II.7)}\end{aligned}$$

for almost all $\mathbf{x} \in \mathcal{X}^{\mathbb{N}}$ in measure $\mathbf{P}(\cdot|\mathbf{y})$, and for almost all $\mathbf{y} \in \mathcal{Y}^{\mathbb{N}}$ in measure $\mathbf{Q}(\cdot|\mathbf{x})$. Note that $p_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1}, \mathcal{Y}_{0,n-1})$ and $q_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1}, \mathcal{X}_n)$ are stochastic kernels (see Definition VI.15, Appendix VI-A) determined from $\mathbf{P}(\cdot|\cdot)$ and $\mathbf{Q}(\cdot|\cdot)$, respectively, (e.g., they are related via (II.1) and (II.4), respectively).

Consequently, the joint distribution of the RV's $\{(X_i, Y_i) : i \in \mathbb{N}\}$ is defined by

$$\begin{aligned}\mathbb{P}\big\{X_0 \in A_0, Y_0 \in B_0, \ldots, X_n \in A_n, Y_n \in B_n\big\} = \int_{A_0} p_0(dx_0) \int_{B_0} q_0(dy_0; x_0) \ldots \\ \int_{A_n} p_n(dx_n; x^{n-1}, y^{n-1}) \int_{B_n} q_n(dy_n; y^{n-1}, x^n).\end{aligned} \qquad \text{(II.8)}$$

Hence, given the two Polish spaces $\mathcal{X}^{\mathbb{N}}$ and $\mathcal{Y}^{\mathbb{N}}$, for any $\mathbf{P}(\cdot|\cdot)$ and $\mathbf{Q}(\cdot|\cdot)$ satisfying the consistency conditions **C1**, **C2**, respectively, there exist a probability space and a sequence of RV's $\{(X_i, Y_i) : i \in \mathbb{N}\}$ defined on it, whose joint probability distribution is uniquely defined

by (II.8), via $\mathbf{P}(\cdot|\cdot)$ and $\mathbf{Q}(\cdot|\cdot)$.

The following remark summarizes the previous discussion on the two equivalent definitions of forward and feedback channels using the definition of stochastic kernels $\big($regular conditional distributions and stochastic kernels are equivalent notions, see Appendix VI-A, Definition VI.13$\big)$.

Remark II.1.

Suppose $\{\mathcal{X}_n : n \in \mathbb{N}\}$, $\{\mathcal{Y}_n : n \in \mathbb{N}\}$, *are complete, separable metric spaces (Polish spaces) and* $\{\mathcal{B}(\mathcal{X}_n) : n \in \mathbb{N}\}$, $\{\mathcal{B}(\mathcal{Y}_n) : n \in \mathbb{N}\}$ *are respectively, the* $\sigma-$*algebras of Borel sets.*

Then

1) *The collection of stochastic kernels*[2] $\{p_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1}) : n \in \mathbb{N}\}$ *uniquely define a family of probability measures on* $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ *via (II.1).*
2) *For any family of probability measures* $\mathbf{P}(\cdot|\mathbf{y})$ *on* $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ *satisfying consistency condition* **C1** *there exists a collection of stochastic kernels* $\{p_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1}) : n \in \mathbb{N}\}$ *connected to* $\mathbf{P}(\cdot|\cdot)$ *via (II.1).*
3) *The collection of stochastic kernels* $\{q_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n}) : n \in \mathbb{N}\}$ *uniquely define a family of probability measures on* $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ *via (II.4).*
4) *For any family of probability measures* $\mathbf{Q}(\cdot|\mathbf{x})$ *on* $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ *satisfying consistency condition* **C2** *there exists a collection of stochastic kernels* $\{q_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n}) : n \in \mathbb{N}\}$ *connected to* $\mathbf{Q}(\cdot|\cdot)$ *via (II.4).*

The point to be made here is that directed information as defined by (I.1)-(I.3) can be expressed via the equivalent definitions of Remark II.1, 2) and 4) rather than 1) and 3). Via this equivalent definition of directed information, the functional and topological properties of mutual information $I(X^n;Y^n)$ can be extended to directed information on general abstract spaces. Throughout the rest of the paper it is assumed that the conditions of Remark II.1 are satisfied. Indeed, according to Remark II.1, all that is required is Polish spaces, which is assumed throughout the rest of the paper. However, if the spaces $\{(\mathcal{X}_n, \mathcal{Y}_n) : n \in \mathbb{N}\}$ are not complete, countably additivity of the family of probability measures $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$ does not fail. This is because these spaces can be homeomorphically embedded as Borel subsets in complete separable metric spaces $\{(\tilde{\mathcal{X}}_n, \tilde{\mathcal{Y}}_n) : n \in \mathbb{N}\}$, so that Kolmogorov's extension theorem can be utilized [22].

[2] $\mathcal{Q}(\mathcal{X};\mathcal{Y} \times \mathcal{Z})$ denotes the set of stochastic kernels on $\mathcal{X}$ given $\mathcal{Y} \times \mathcal{Z}$.

III. Properties of Directed Information

In this section, we define the feedforward information $I(X^n \to Y^n)$ and the feedback information $I(X^n \leftarrow Y^n)$ on abstract spaces (Polish spaces), via the Kullback-Leibler distance (or relative entropy), using the basic family of measures $\mathbf{P}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$, and $\mathbf{Q}(\cdot|\mathbf{x})$ on $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$, which satisfy consistency condition $\mathbf{C1}$ and $\mathbf{C2}$, respectively, and Radon-Nikodym Derivatives (RNDs). Once this is established, then following Pinsker [29] it will become obvious that directed information permits a representation as a supremum of relative entropy between two distributions, where the supremum is taken over all measurable partitions on a given $\sigma-$ algebra of subsets of a set $\mathcal{Z}$. Further, in a subsequent subsection, we use the definition of directed information in terms of $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$, to derive several of its properties, such as, convexity, concavity, lower semicontinuity, with respect to these two families of measures.

To present the precise expression for the directed information, following Csiszár [20], we first introduce the measures of interest constructed from the basic consistent families of conditional distributions. Let $\mathcal{M}_1(\Sigma)$ denote the set of probability measures on the measurable space (Σ, $\mathcal{B}(\Sigma)$). Introduce the following notation.

$$\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}}) \stackrel{\triangle}{=} \Big\{ \mathbf{P}(\cdot|\mathbf{x}) \in \mathcal{M}_1(\mathcal{X}^{\mathbb{N}}) : \mathbf{P}(\cdot|\mathbf{x}) \text{ are regular probability measures satisfying consistency condition } \mathbf{C1} \Big\}$$

$$\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}}) \stackrel{\triangle}{=} \Big\{ \mathbf{Q}(\cdot|\mathbf{y}) \in \mathcal{M}_1(\mathcal{Y}^{\mathbb{N}}) : \mathbf{Q}(\cdot|\mathbf{y}) \text{ are regular probability measures satisfying consistency condition } \mathbf{C2} \Big\}.$$

The projection of $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$ to finite number of coordinates is denoted by $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$, respectively.

Next, we define the distributions of interest. Given any conditional distributions $\mathbf{P}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$, utilizing the construction of Section II, we can define uniquely $\{p_n(\cdot;\cdot,\cdot) : n \in \mathbb{N}\}$ and $\{q_n(\cdot;\cdot,\cdot) : n \in \mathbb{N}\}$, $\big($see (II.6), (II.7)$\big)$ and the following distributions.

P1: The joint distribution on $\mathcal{X}^{\mathbb{N}} \times \mathcal{Y}^{\mathbb{N}}$ of the basic sequence $\{X_n, Y_n : n \in \mathbb{N}\}$ constructed from $\mathbf{P}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$, defined uniquely for $A_i \in \mathcal{B}(\mathcal{X}_i)$,

$B_i \in \mathcal{B}(\mathcal{Y}_i)$, $i \in \mathbb{N}^n$, by

$$(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(\times_{i=0}^n (A_i \times B_i)) \stackrel{\triangle}{=} \mathbb{P}\Big\{X_0 \in A_0, Y_0 \in B_0, \ldots, X_n \in A_n, Y_n \in B_n\Big\}$$
$$= \int_{A_0} p_0(dx_0) \int_{B_0} q_0(dy_0; x_0) \ldots \int_{A_n} p_n(dx_n; x^{n-1}, y^{n-1}) \int_{B_n} q_n(dy_n; y^{n-1}, x^n). \quad \text{(III.1)}$$

Formally, (III.1) is written as $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n)$ or $\overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \otimes \overrightarrow{Q}_{0,n}(dy^n|x^n)$.

P2: The marginal distributions on $\mathcal{X}^{\mathbb{N}}$ of the sequence $\{X_n : n \in \mathbb{N}\}$ constructed from $\mathbf{P}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$, defined uniquely by[3]

$$\mu_{0,n}(\times_{i=0}^n A_i) \stackrel{\triangle}{=} \mathbb{P}\Big\{X_0 \in A_0, Y_0 \in \mathcal{Y}_0, \ldots, X_n \in A_n, Y_n \in \mathcal{Y}_n\Big\}, \ A_i \in \mathcal{B}(\mathcal{X}_i), \ i \in \mathbb{N}^n \quad \text{(III.2)}$$
$$= (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(\times_{i=0}^n (A_i \times \mathcal{Y}_i))$$
$$= \int_{A_0} p_0(dx_0) \int_{\mathcal{Y}_0} q_0(dy_0; x_0) \ldots \int_{A_n} p_n(dx_n; x^{n-1}, y^{n-1}) \int_{\mathcal{Y}_n} q_n(dy_n; y^{n-1}, x^n). \quad \text{(III.3)}$$

Formally, (III.2) is written as $\mu_{0,n}(dx^n) = (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, \mathcal{Y}_{0,n})$, and by Bayes' rule $\mu_{0,n}(dx^n) = \otimes_{i=0}^n \mu_{i|i-1}(dx_i; x^{i-1})$.

P3: The marginal distributions on $\mathcal{Y}^{\mathbb{N}}$ of the sequence $\{Y_n : n \in \mathbb{N}\}$ constructed from $\mathbf{P}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$, defined uniquely by

$$\nu_{0,n}(\times_{i=0}^n B_i) \stackrel{\triangle}{=} \mathbb{P}\Big\{X_0 \in \mathcal{X}_0, Y_0 \in B_0, \ldots, X_n \in \mathcal{X}_n, Y_n \in B_n\Big\}, \ B_i \in \mathcal{B}(\mathcal{Y}_i), \ i \in \mathbb{N}^n \quad \text{(III.4)}$$
$$= (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(\times_{i=0}^n (\mathcal{X}_i \times B_i))$$
$$= \int_{\mathcal{X}_0} p_0(dx_0) \int_{B_0} q_0(dy_0; x_0) \ldots \int_{\mathcal{X}_n} p_n(dx_n; x^{n-1}, y^{n-1}) \int_{B_n} q_n(dy_n; y^{n-1}, x^n). \quad \text{(III.5)}$$

Formally, (III.4) is written as $\nu_{0,n}(dy^n) = (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n}, dy^n)$, and by Bayes' rule $\nu_{0,n}(dy^n) = \otimes_{i=0}^n \nu_{i|i-1}(dy_i; y^{i-1})$.

P4: The measure $\overrightarrow{\Pi}_{0,n} : \mathcal{B}(\mathcal{X}_{0,n}) \odot \mathcal{B}(\mathcal{Y}_{0,n}) \mapsto [0,1]$ constructed from $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$

[3]Actually $\mu \equiv \mu^{\mathbf{P} \otimes \mathbf{Q}}$ but we omit the superscript throughout the paper.

and $\nu_{0,n}(dy^n) = (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n}, dy^n) \in \mathcal{M}_1(\mathcal{Y}_{0,n})$ of (III.4), defined uniquely by

$$\begin{aligned}\overrightarrow{\Pi}_{0,n}(\times_{i=0}^n (A_i \times B_i)) &\stackrel{\triangle}{=} (\overleftarrow{P}_{0,n} \otimes \nu_{0,n})(\times_{i=0}^n (A_i \times B_i)),\ A_i \in \mathcal{B}(\mathcal{X}_i),\ B_i \in \mathcal{B}(\mathcal{Y}_i),\ i \in \mathbb{N}^n \\ &= \int_{A_0} p_0(dx_0) \int_{B_0} \nu_0(dy_0) \int_{A_1} p_1(dx_1; x_0, y_0) \int_{B_1} \nu_{1|0}(dy_1; y_0) \dots \\ &\dots \int_{A_n} p_n(dx_n; x^{n-1}, y^{n-1}) \int_{B_n} \nu_{n|n-1}(dy_n; y^{n-1}). \end{aligned} \tag{III.6}$$

Formally, (III.6) is written as $\overrightarrow{\Pi}_{0,n}(dx^n, dy^n) = \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \otimes \nu_{0,n}(dy^n)$.

P5: The measure $\overleftarrow{\Pi}_{0,n} : \mathcal{B}(\mathcal{Y}_{0,n}) \odot \mathcal{B}(\mathcal{X}_{0,n}) \mapsto [0,1]$ constructed from $\overrightarrow{Q}_{0,n}(dy^n|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ and $\mu_{0,n}(dx^n) = (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, \mathcal{Y}_{0,n}) \in \mathcal{M}_1(\mathcal{X}_{0,n})$ of (III.3), defined uniquely by

$$\begin{aligned}\overleftarrow{\Pi}_{0,n}(\times_{i=0}^n (A_i \times B_i)) &\stackrel{\triangle}{=} (\mu_{0,n} \otimes \overrightarrow{Q}_{0,n})(\times_{i=0}^n (A_i \times B_i)),\ A_i \in \mathcal{B}(\mathcal{X}_i),\ B_i \in \mathcal{B}(\mathcal{Y}_i),\ i \in \mathbb{N}^n \\ &= \int_{A_0} \mu_0(dx_0) \int_{B_0} q_0(dy_0; x_0) \int_{A_1} \mu_{1|0}(dx_1; x_0) \int_{B_1} q_1(dy_1; y_0, x_0) \dots \\ &\dots \int_{A_n} \mu_{n|n-1}(dx_n; x^{n-1}) \int_{B_n} q_n(dy_n; y^{n-1}, x^n). \end{aligned} \tag{III.7}$$

Formally, (III.7) is written as $\overleftarrow{\Pi}_{0,n}(dx^n, dy^n) = \mu_{0,n}(dx^n) \otimes \overrightarrow{Q}_{0,n}(dy^n|x^n)$.

From the above definitions, an alternative way to construct the conditional distributions $\nu_{n|n-1}(\cdot;\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1})$ and $\mu_{n|n-1}(\cdot;\cdot) \in \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1})$ is as follows. Let $A^{(n)} = \{\mathbf{x} : x_n \in A\}$, $A \in \mathcal{B}(\mathcal{X}_n)$, $B^{(n)} = \{\mathbf{y} : y_n \in B\}$, $B \in \mathcal{B}(\mathcal{Y}_n)$, and let $\overrightarrow{\Pi}_{0,n}(A^{(n)}, B^{(n)}|\mathcal{B}(\mathcal{X}_{0,n-1}) \odot \mathcal{B}(\mathcal{Y}_{0,n-1}))$ denote the joint conditional probability of $A^{(n)} \times B^{(n)}$ with respect to $\mathcal{B}(\mathcal{X}_{0,n-1}) \odot \mathcal{B}(\mathcal{Y}_{0,n-1})$ calculated on the probability space $\left(\mathcal{X}^{\mathbb{N}} \otimes \mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}) \odot \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \overrightarrow{\Pi}_{0,n}(\cdot)\right)$. Then for $A \in \mathcal{B}(\mathcal{X}_n)$, $B \in \mathcal{B}(\mathcal{Y}_n)$ we obtain

$$\overrightarrow{\Pi}_{0,n}(A^{(n)}, B^{(n)}|\mathcal{B}(\mathcal{X}_{0,n-1}) \odot \mathcal{B}(\mathcal{Y}_{0,n-1})) = p_n(A; x^{n-1}, y^{n-1}) \times \nu_{n|n-1}(B; y^{n-1}). \tag{III.8}$$

Hence, $\nu_{n|n-1}(\cdot;\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1})$ is given by $\nu_{n|n-1}(dy_n; y^{n-1}) = \int_{\mathcal{X}_n} \overrightarrow{\Pi}_{0,n}(dx_n, dy_n; x^{n-1}, y^{n-1})$, from which $\nu_{0,n}(dy^n) \in \mathcal{M}_1(\mathcal{Y}_{0,n})$ is also obtained. Similarly, let $\overleftarrow{\Pi}_{0,n}(A^{(n)}, B^{(n)}|\mathcal{B}(\mathcal{Y}_{0,n-1}) \odot \mathcal{B}(\mathcal{X}_{0,n-1}))$ denote the joint conditional probability of $A^{(n)} \times B^{(n)}$ with respect to $\mathcal{B}(\mathcal{Y}_{0,n-1}) \odot \mathcal{B}(\mathcal{X}_{0,n-1})$ calculated on the probability space $\left(\mathcal{Y}^{\mathbb{N}} \times \mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}) \odot \mathcal{B}(\mathcal{X}^{\mathbb{N}}), \overleftarrow{\Pi}_{0,n}(\cdot)\right)$. Then for $B \in \mathcal{B}(\mathcal{Y}_n)$,

$$\overleftarrow{\Pi}_{0,n}(A^{(n)}, B^{(n)}|\mathcal{B}(\mathcal{X}_{0,n-1}) \odot \mathcal{B}(\mathcal{Y}_{0,n-1})) = \int_{A_n} q_n(B; y^{n-1}, x^n) \otimes \mu_{n|n-1}(dx_n; x^{n-1}) \tag{III.9}$$

from which $\mu_{n|n-1}(\cdot;\cdot) \in \mathcal{Q}(\mathcal{X}_n;\mathcal{X}_{0,n-1})$ and $\mu_{0,n}(dx^n) \in \mathcal{M}_1(\mathcal{X}_{0,n})$ are obtained. Similarly, from (III.6) and (III.8) we can obtain any of the individual kernels $p_n(\cdot;\cdot,\cdot)$ and $q_n(\cdot;\cdot,\cdot)$ appearing in their right hand side by proper conditional expectations.

Using the first definition of basic processes, that is, given a collection of stochastic kernels $\{p_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{X}_n;\mathcal{X}_{0,n-1}\times\mathcal{Y}_{0,n-1}) : n \in \mathbb{N}\}$ and $\{q_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{Y}_n;\mathcal{Y}_{0,n-1}\times\mathcal{X}_{0,n}) : n \in \mathbb{N}\}$, the joint distribution, as well as the conditional distributions are defined via $\mathbf{P1}-\mathbf{P5}$. Consequently, it is well-known that directed information is defined via relative entropy as follows [8] (see Appendix VI-A, pp. 47-48).

$$I(X^n \to Y^n) \stackrel{\triangle}{=} \sum_{i=0}^{n} I(X^i;Y_i|Y^{i-1})$$

$$= \sum_{i=0}^{n} \int_{\mathcal{Y}_{0,i-1}} \int_{\mathcal{X}_{0,i}\times\mathcal{Y}_i} \log\left(\frac{P_{0,i}(dx^i,dy_i;y^{i-1})}{P_{0,i}(dx^i;y^{i-1})\times\nu_{i|i-1}(dy_i;y^{i-1})}\right) P_{0,i}(dx^i,dy_i;y^{i-1})P_{0,i-1}(dy^{i-1}) \tag{III.10}$$

$$= \sum_{i=0}^{n} \int_{\mathcal{X}_{0,i}\times\mathcal{Y}_{0,i-1}} \mathbb{D}(q_i(\cdot;y^{i-1},x^i)||\nu_{i|i-1}(\cdot;y^{i-1}))p_i(dx_i;x^{i-1},y^{i-1})$$
$$\otimes_{j=0}^{i-1}\left(q_j(dy_j;y^{j-1},x^j)\otimes p_j(dx_j;x^{j-1},y^{j-1})\right) \tag{III.11}$$

$$\equiv \mathbb{I}_{X^n\to Y^n}(p_i(\cdot;\cdot,\cdot),q_i(\cdot;\cdot,\cdot): \ i=0,1,\ldots,n). \tag{III.12}$$

The right hand side in (III.10) follows from the definition of conditional mutual information, while we use the notation $\mathbb{I}_{X^n\to Y^n}(p_i(\cdot;\cdot,\cdot),q_i(\cdot;\cdot,\cdot): \ i=0,1,\ldots,n)$ to denote that $I(X^n \to Y^n)$ is a functional of $\{p_i(\cdot;\cdot,\cdot),q_i(\cdot;\cdot,\cdot): \ i=0,1,\ldots,n\}$. Additional comments regarding (III.10)-(III.12) are given in Appendix VII-A (pp. 48-50).

A. Directed Information Functional of Consistent Conditional Distributions

Now we consider the second definition of basic process introduced in Section II. Given any family of measures $\mathbf{P}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ the measures under $\mathbf{P1}-\mathbf{P5}$ are constructed. Next, we define directed information via relative entropy as done for mutual information [20]. By Lemma VI.14, $\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n} << \overleftarrow{P}_{0,n}\otimes\nu_{0,n}$ if and only if $\overrightarrow{Q}_{0,n}(\cdot|x^n) << \nu_{0,n}(\cdot)$ for $\overleftarrow{P}_{0,n}-$almost all $x^n \in \mathcal{X}_{0,n}$. Utilizing the Radon-Nikodym derivative (RND) $\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}(x^n,y^n)$, define the relative entropy of $\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}$ with respect to $\overrightarrow{\Pi}_{0,n}$ as

follows.

$$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n}) \stackrel{\triangle}{=} \mathbb{D}(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}||\overrightarrow{\Pi}_{0,n})$$

$$= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}\Big)(x^n,y^n)d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n) \tag{III.13}$$

$$= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n). \tag{III.14}$$

Note that (III.14) is obtained by utilizing the fact that if $\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n} << \overleftarrow{P}_{0,n}\otimes\nu_{0,n}$ then the RND $\frac{(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}(x^n,y^n)$ represents a version of $\frac{\overrightarrow{Q}_{0,n}(\cdot|x^n)}{\nu_{0,n}(\cdot)}(y^n)$, $\overleftarrow{P}_{0,n}-a.s$ for all $x^n\in\mathcal{X}_{0,n}$. On the other hand, using Lemma VI.14, $\overrightarrow{Q}_{0,n}(\cdot|x^n)\ll\nu_{0,n}(\cdot)$, $\overleftarrow{P}_{0,n}-$almost $x^n\in\mathcal{X}_{0,n}$, and by Radon-Nikodym theorem, there exists a version of the RND $\bar{\xi}_{0,n}(x^n,y^n)\stackrel{\triangle}{=}\frac{\overrightarrow{Q}_{0,n}(\cdot|x^n)}{\nu_{0,n}(\cdot)}(y^n)$ which is a non-negative measurable function of $(x^n,y^n)\in\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}$. Hence another version of $\bar{\xi}_{0,n}(\cdot,\cdot)$ is $\bar{\xi}_{0,n}(x^n,y^n)=\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}(x^n,y^n)$. We use the previous notation $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})$ to illustrate that $\mathbb{D}(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}||\overrightarrow{\Pi}_{0,n})$ is a functional of $\big\{\overleftarrow{P}_{0,n}(\cdot|y^{n-1}),\overrightarrow{Q}_{0,n}(\cdot|x^n)\big\}\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\times\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$.

In the next Remark we summarize the equivalent definitions of directed information based on the two equivalent definitions of channels, that is, the one based on (III.11), (III.12), and the one based on (III.13), (III.14).

Remark III.1.

Let $\mathbf{P}(\cdot|\cdot)\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ *and* $\mathbf{Q}(\cdot|\cdot)\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$. *By repeated application of Lemma VI.14, and the chain rule of relative entropy, Theorem VI.19 of Appendix VI-A, directed information*

admits the following equivalent definitions.

$$I(X^n \to Y^n) \stackrel{\triangle}{=} \sum_{i=0}^{n} I(X^i; Y_i|Y^{i-1}) \tag{III.15}$$

$$= \sum_{i=0}^{n} \int_{\mathcal{X}_{0,i}\times\mathcal{Y}_{0,i-1}} \left(\int_{\mathcal{Y}_i} \mathbb{D}(q_i(\cdot; y^{i-1}, x^i)||\nu_{i|i-1}(\cdot; y^{i-1})) \right) p_i(dx_i; x^{i-1}, y^{i-1})$$

$$\otimes_{j=0}^{i-1} \Big(q_j(dy_j; y^{j-1}, x^j) \otimes p_j(dx^j; x^{j-1}, y^{j-1}) \Big) \tag{III.16}$$

$$= \mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n}||\overrightarrow{\Pi}_{0,n}) \tag{III.17}$$

$$= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log \Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)} \Big) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) \tag{III.18}$$

$$\equiv \mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}). \tag{III.19}$$

Clearly, (III.19) is valid even when $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n)$ is singular with respect to $(\overleftarrow{P}_{0,n} \otimes \nu_{0,n})(dx^n, dy^n)$, in which case its value is $+\infty$. The point to be made here is that we will show the convexity, concavity, lower semicontinuity properties of directed information $I(X^n \to Y^n)$, using the functional $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$, with respect to the family of measures $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$. We will also use the directed information definition $\mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n}||\overrightarrow{\Pi}_{0,n})$, as a functional of $\{\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}\}$ to show lower semicontinuity, convexity and concavity properties. These functional and topological properties are important in showing existence of extremum problems, such as, maximizing directed information over channel input distributions $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ satisfying power constraints, for capacity calculations, and minimizing directed information over $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ satisfying fidelity constraints for sequential and nonanticipative rate distortion functions.

Remark III.2.

Note that one may investigate directed information $\mathbb{I}_{X^n\to Y^n}(p_i(\cdot;\cdot,\cdot), q_i(\cdot;\cdot,\cdot) : \ i = 0, 1, \ldots, n)$ *as a functional on the space of vector measures* $\{p_0, p_1, \ldots, p_n\} \in \times_{i=0}^{n} \mathcal{Q}(\mathcal{X}_i; \mathcal{X}_{0,i-1} \times \mathcal{Y}_{0,i-1})$ *and* $\{q_0, q_1, \ldots, q_n\} \in \times_{i=0}^{n} \mathcal{Q}(\mathcal{Y}_i; \mathcal{Y}_{0,i-1} \times \mathcal{X}_{0,i})$*. However, it is not clear to us whether this formulation is appropriate to obtain the results R1)–R7).*

Feedback Information.

Similar to the above discussion we can also define feedback information as follows. It can be

easily verified that $\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1} << \overrightarrow{Q}_{0,n-1} \times \mu_{0,n}$ if and only if $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) << \mu_{0,n}(\cdot)$ for $\nu_{0,n-1}-$almost all $y^{n-1} \in \mathcal{Y}_{0,n-1}$. Moreover, if $\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1} << \overrightarrow{Q}_{0,n-1} \otimes \mu_{0,n}$ then the RND $\frac{d(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1})}{d(\overrightarrow{Q}_{0,n-1} \otimes \mu_{0,n})}(x^n, y^{n-1})$ represents a version of $\frac{d\overleftarrow{P}_{0,n}(\cdot|y^{n-1})}{d\mu_{0,n}(\cdot)}(x^n)$, $\nu_{0,n-1} - a.s$ for all $y^{n-1} \in \mathcal{Y}_{0,n-1}$. Hence, when $\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1} << \overrightarrow{Q}_{0,n-1} \otimes \mu_{0,n}$, then

$$
\begin{aligned}
I(X^n \leftarrow Y^n) &\stackrel{\triangle}{=} \sum_{i=0}^{n} I(Y^{i-1}; X_i | X^{i-1}) \\
&= \sum_{i=0}^{n} \int_{\mathcal{X}_{0,i-1} \times \mathcal{Y}_{0,i-1}} \left(\int_{\mathcal{X}_i} \mathbb{D}(p_i(\cdot; x^{i-1}, y^{i-1}) || \mu_{i|i-1}(\cdot; x^{i-1})) \right) \\
&\qquad \otimes_{j=0}^{i-1} \Big(p_j(dx^j; x^{j-1}, y^{j-1}) \otimes q_j(dy_j; y^{j-1}, x^j) \Big) \\
&= \mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1} || \overleftarrow{\Pi}_{0,n}) \\
&\stackrel{\triangle}{=} \int_{\mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n}} \log \Big(\frac{d(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1})}{d(\overrightarrow{Q}_{0,n-1} \otimes \mu_{0,n})} \Big)(x^n, y^{n-1}) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1})(dx^n, dy^{n-1}) \qquad \text{(III.20)} \\
&= \int_{\mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n}} \log \Big(\frac{\overleftarrow{P}_{0,n}(dx^n|y^{n-1})}{\mu_{0,n}(dx^n)} \Big) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n-1})(dx^n, dy^{n-1}) \qquad \text{(III.21)} \\
&\equiv \mathbb{I}_{X^n \leftarrow Y^n}(\overrightarrow{Q}_{0,n-1}, \overleftarrow{P}_{0,n}). \qquad \text{(III.22)}
\end{aligned}
$$

Note that (III.22) states that directed information $I(X^n \leftarrow Y^n)$ is expressed as a functional of $\{\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n-1}\}$ and it is denoted by $\mathbb{I}_{X^n \leftarrow Y^n}(\overrightarrow{Q}_{0,n-1}, \overleftarrow{P}_{0,n})$. We will not pursue the properties of (III.22) since these follow along the same line of reasoning as those of $I(X^n \rightarrow Y^n)$.

We also point out that directed information can be equivalently defined as a supremum over appropriate partition of measurable spaces with respect to the families $\mathbf{P}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$. This construction is elaborated in Appendix VII-A.

We are now ready to investigate the functional properties of directed information, as a functional of the space of input conditional distributions $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$, and the space of channel conditional distributions $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$, using the relative entropy definition of directed information, $I(X^n \rightarrow Y^n) = \mathbb{I}_{X^n \rightarrow Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$, as a function of $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$. Recall that for mutual information defined via relative entropy, analogous properties, such as, convexity, concavity, lower semicontinuity etc, are well known

[20], [27], [30]. We shall derive similar properties for the functional $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$.

B. Convexity and Concavity of Directed Information

First, we show that the family of regular conditional distributions $\mathbf{P}(\cdot|\mathbf{y})$ and $\mathbf{Q}(\cdot|\mathbf{x})$ satisfying consistency conditions **C1** and **C2**, are convex, and then we show convexity of directed information with respect to $\mathbf{Q}(\cdot|\mathbf{x})$ and concavity with respect to $\mathbf{P}(\cdot|\mathbf{y})$. We recall the definition of convexity of the set of regular conditional measures given in Definition VI.17 (see Appendix VI-A, p. 47). Let $\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big)$ denote the set of probability measures on $(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$, which are absolutely continuous with respect to a fixed reference probability measure $\bar{\mathbf{P}}$, and have restrictions on $\mathcal{B}(\mathcal{Y}^{\mathbb{N}})$ which are equivalent (mutually absolutely continuous) with respect to $\bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}$. Hence, for any $\mathbf{P}\in\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big)$, by definition, $\mathbf{P}\ll\bar{\mathbf{P}}$ and $\mathbf{P}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}\ll\bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}$, $\bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}\ll\mathbf{P}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}$ denoted by $\mathbf{P}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}\sim\bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{Y}^{\mathbb{N}})}$.
Introduce the set of all regular conditional probability measures

$$\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}^{\mathbb{N}})\Big)(\mathbf{y})$$
$$\stackrel{\triangle}{=}\Big\{\mathbf{P}(\cdot|\mathbf{y}) : \mathbf{P}\in\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big)\Big\}. \quad \text{(III.23)}$$

Define the subset of the above set consisting of all conditional distributions which satisfy consistency condition $\mathbf{C1}$ as follows.

$$\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})\stackrel{\triangle}{=}\Big\{\mathbf{P}(\cdot|\mathbf{y})\in\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}^{\mathbb{N}})\Big)(\mathbf{y}) :$$
$$\mathbf{P}(\cdot|\mathbf{y}) \text{ satisfies consistency condition } \mathbf{C1}\Big\}.$$

The projection to finite coordinates is defined by

$$\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\stackrel{\triangle}{=}\Big\{\mathbf{P}(\cdot|\mathbf{y})\in\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\Big)(y^{n-1}) :$$
$$\mathbf{P}(\cdot|\mathbf{y}) \text{ satisfies consistency condition } \mathbf{C1}\Big\}. \quad \text{(III.24)}$$

Similarly, define the set of measures $\mathbf{Q}(\cdot|\mathbf{x})$ which satisfy consistency condition $\mathbf{C2}$ by

$$\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})\stackrel{\triangle}{=}\Big\{\mathbf{Q}(\cdot|\mathbf{x})\in\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{X}^{\mathbb{N}})\Big)(\mathbf{x}) :$$
$$\mathbf{Q}(\cdot|\mathbf{x}) \text{ satisfies consistency condition } \mathbf{C2}\Big\}, \quad \text{(III.25)}$$

and its projection to finite coordinates by

$$\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n}) \triangleq \Big\{\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{X}_{0,n})\Big)(x^n) : \mathbf{Q}(\cdot|\mathbf{x}) \text{ satisfies consistency condition } \mathbf{C2}\Big\}. \quad \text{(III.26)}$$

Next, we show convexity of the sets $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$.

Theorem III.3.

Let $\{\mathcal{X}_n : n\in\mathbb{N}\}$, $\{\mathcal{Y}_n : n\in\mathbb{N}\}$ be Polish spaces with $\mathcal{B}(\mathcal{X}_n)$, $\mathcal{B}(\mathcal{Y}_n)$, respectively, the $\sigma-$algebras of Borel sets. Then the sets $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ are convex subsets of the sets of regular conditional distributions $\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\Big)(y^{n-1})$ and $\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{X}_{0,n})\Big)(x^n)$, respectively.

Proof. The a.s.-convexity of the sets $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ will be shown using Definition VI.17. Since the methodology is similar for both sets, only the derivation of $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ will be given. It is well-known that the sets $\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\Big)(y^{n-1})$ and $\mathcal{M}_1\Big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{X}_{0,n})\Big)(x^n)$ are convex. By the definition of a.s.-convexity of the set of regular conditional distributions (Definition VI.17), $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ is convex (almost surely) if for a given $\mathbf{P}^1(\cdot|\mathbf{y}), \mathbf{P}^2(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$, and a given $\lambda\in(0,1)$, there exists a probability measure $\tilde{\mathbf{P}}$ on $(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$, whose regular conditional measure $\tilde{\mathbf{P}}(\cdot|\mathbf{y}) \in \mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)(y^{n-1})$ is a convex combination $\tilde{\mathbf{P}}(\cdot|\mathbf{y}) = \lambda\mathbf{P}^1(\cdot|\mathbf{y}) + (1-\lambda)\mathbf{P}^2(\cdot|\mathbf{y})$, $\bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{Y}_{0,n-1})} - a.s.$, and consistency condition $\mathbf{C1}$ holds. This property is denoted by $\lambda\mathbf{P}^1(\cdot|\mathbf{y}) + (1-\lambda)\mathbf{P}^2(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. By [31] the set of regular conditional distributions $\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)(y^{n-1})$ is a convex set. Also, $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1}) \subset \mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)(y^{n-1})$, and hence if $\mathbf{P}^1(\cdot|\mathbf{y}), \mathbf{P}^2(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ then they are included in $\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)(y^{n-1})$, and hence they satisfy the property

$$\lambda\mathbf{P}^1(\cdot|\mathbf{y}) + (1-\lambda)\mathbf{P}^2(\cdot|\mathbf{y}) \in \mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)(y^{n-1}), \quad \forall\lambda\in(0,1).$$

To complete the derivation it is sufficient to show that the consistency condition $\mathbf{C1}$ holds. Let $A^{(n)} = \{\mathbf{x} : x_n\in A\}$, $A\in\mathcal{B}(\mathcal{X}_n)$. In addition, let $\mathbf{P}(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1}))$ denote the conditional probability of $A^{(n)}$ with respect to $\mathcal{B}(\mathcal{X}_{0,n-1})$ calculated on the probability space $\big(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}), \mathbf{P}(\cdot|\mathbf{y})\big)$. From the definition of regular conditional probability measures on $\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot$

$\mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)$ it follows that

$$\begin{aligned}\tilde{\mathbf{P}}(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1})) &= \lambda\mathbf{P}^1(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1})) + (1-\lambda)\mathbf{P}^2(A^{(n)}|\mathbf{y}|\mathcal{B}(\mathcal{X}_{0,n-1})) - a.s. \\ &= \lambda p_n^1(dx_n; x^{n-1}, y^{n-1}) + (1-\lambda)p_n^2(dx_n; x^{n-1}, y^{n-1}) - a.s.\end{aligned}$$

where $p_n^1(\cdot;\cdot,\cdot)$ and $p_n^2(\cdot;\cdot,\cdot)$ are regular conditional distributions. Since convex combination of regular conditional distributions is also a regular conditional distribution, then $\lambda\mathbf{P}^1(\cdot|\mathbf{y}) + (1-\lambda)\mathbf{P}^2(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$, e.g, the consistency condition **C1** holds, and the derivation is complete. The derivation of a.s.-convexity of the set $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ is done similarly. □

Since $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ are convex, then we proceed further to show that directed information $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$, as a functional of $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ for a fixed $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ is concave, and as a functional of $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ for a fixed $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ is convex. These results are shown in the next theorem.

Theorem III.4.

Let $\{\mathcal{X}_n : n \in \mathbb{N}\}$, $\{\mathcal{Y}_n : n \in \mathbb{N}\}$ be Polish spaces with $\mathcal{B}(\mathcal{X}_n)$, $\mathcal{B}(\mathcal{Y}_n)$, respectively, the $\sigma-$algebras of Borel sets. Consider the directed information functional $I(X^n \to Y^n) = \mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$, $\mathbb{I}_{X^n\to Y^n} : \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1}) \times \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n}) \mapsto [0,\infty]$ defined by (III.19).

Then the following hold.

1) *$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ is a convex functional of $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ for a fixed $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$.*
2) *$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ is a concave functional of $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ for a fixed $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$.*
3) *$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \cdot)$ is a strictly convex functional on the set $\big\{\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n}) : \mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}) < \infty\big\}$ for a fixed $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$.*

Proof. By Theorem III.3, the sets $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ are convex. Therefore, to show parts 1), 2), 3) we utilize the consistency of the two families of conditional distributions and we apply the log-sum formulae. The complete derivation is given in Appendix VII-B. □

Theorem III.4 is analogous to similar results for mutual information $I(X^n;Y^n) \equiv \mathbb{I}_{X^n;Y^n}(P_{X^n}, P_{Y^n|X^n})$, expressed as a functional of input distribution $P_{X^n}(dx^n) \in \mathcal{M}_1(\mathcal{X}_{0,n})$ and the channel $P_{Y^n|X^n}(dy^n|x^n) \in$

$\mathcal{Q}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$, which is a convex (respectively concave) functional of $P_{Y^n|X^n}(dy^n|x^n) \in \mathcal{Q}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ $\big($respectively $P_{X^n}(dx^n) \in \mathcal{M}_1(\mathcal{X}_{0,n})\big)$ for a fixed $P_{X^n}(dx^n) \in \mathcal{M}_1(\mathcal{X}_{0,n})$ $\big($respectively $P_{Y^n|X^n}(dy^n|x^n) \in \mathcal{Q}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})\big)$. It is important to point out that if one considers the alternative definition of directed information (III.10), (III.12), as a functional of the sequence of input channel distributions, $I(X^n \to Y^n) \equiv \mathbb{I}_{X^n\to Y^n}(p_i(\cdot;\cdot,\cdot), q_i(\cdot;\cdot,\cdot) : \ i = 0,1,\ldots,n)$, then it is not clear to us whether it is possible to establish convexity and concavity with respect to q_i and p_i.

C. Weak Convergence and Compactness of Conditional Distributions

In this section we give general sufficient conditions for weak compactness of the set of measures $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ and $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$, and compactness of the set of joint and marginal measures with respect to the topology of weak convergence of probability measures. These conditions are sufficient to show lower semicontinuity of $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ for fixed $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ $\big($respectively $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})\big)$ with respect to $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ $\big($respectively $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\big)$. These results are the analogous to the lower semicontinuity of mutual information, extensively utilized in information theory and statistics (see [20], [27]).

Before we state the main theorem, we introduce the following notation. Let $BC(\mathcal{X})$ denote the set of bounded, continuous real-valued function f defined on the metric space $(\mathcal{X}, d)$ endowed with the supremum norm $||f|| = \sup_{x\in\mathcal{X}} |f(x)|$. A sequence of probability measures $\{P_n : n \geq 1\} \subset \mathcal{M}_1(\mathcal{X})$ is said to *converge weakly* to $P \in \mathcal{M}_1(\mathcal{X})$ if [32]

$$\lim_{n\to\infty} \int_{\mathcal{X}} f(x) dP_n(x) = \int_{\mathcal{X}} f(x) dP(x), \ \forall f \in BC(\mathcal{X}).$$

Weak convergence of $\{P_n : n \geq 1\}$ to P is denoted by $P_n \stackrel{w}{\longrightarrow} P$. Appendix VI-A summarizes the main results of weak convergence, compactness, tightness, and Prohorov's theorem which we will invoke to derive the results of this section.

Throughout sequences of points in $\mathcal{X}^{\mathbb{N}}$ and $\mathcal{Y}^{\mathbb{N}}$ are denoted by $\mathbf{x}^{(\alpha)} \stackrel{\triangle}{=} \{x_0^{(\alpha)}, x_1^{(\alpha)}, \ldots\} \in \mathcal{X}^{\mathbb{N}}$, $\mathbf{y}^{(\alpha)} \stackrel{\triangle}{=} \{y_0^{(\alpha)}, y_1^{(\alpha)}, \ldots\} \in \mathcal{Y}^{\mathbb{N}}$, $\alpha = 0,1,2,\ldots$ Moreover, $\mathbf{x}^{(\alpha)} \in \mathcal{X}^{\mathbb{N}}$ is said to converge to $\mathbf{x}^{(0)} \in \mathcal{X}^{\mathbb{N}}$ as $\alpha \longrightarrow \infty$, if $x_n^{(\alpha)} \longrightarrow x_n^{(0)}$, for all $n \in \mathbb{N}$. Sequence of points in $\otimes_{i=1}^n \mathcal{X}_i$ and $\otimes_{i=1}^n \mathcal{Y}_i$ are denoted by $x^{n,(\alpha)} \stackrel{\triangle}{=} \{x_0^{(\alpha)}, x_1^{(\alpha)}, \ldots, x_n^{(\alpha)}\}$ and $y^{n,(\alpha)} \stackrel{\triangle}{=} \{y_0^{(\alpha)}, y_1^{(\alpha)}, \ldots, y_n^{(\alpha)}\}$, $\alpha = 0,1,\ldots,$ respectively.

Next, we state the general novel theorem which is also used to show lower semicontinuity of

directed information. The theorem consists of two parts depending on whether $\mathcal{Y}_{0,n}$ is compact and $p_n(\cdot; x^{n-1}, y^{n-1})$ weakly continuous, or $\mathcal{X}_{0,n}$ is compact and $q_n(\cdot; y^{n-1}, x^n)$ is weakly continuous. In applications of information theory either one of them or both maybe required.

Theorem III.5.

Part A. *Let $\mathcal{Y}_{0,n}$ be a compact Polish space and $\mathcal{X}_{0,n}$ a Polish space. Assume $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ satisfy the following condition.*

CA*: For all $g(\cdot) \in BC(\mathcal{X}_{0,n})$, the function*

$$(x^{n-1}, y^{n-1}) \in \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1} \longmapsto \int_{\mathcal{X}_n} g(x) p_n(dx; x^{n-1}, y^{n-1}) \in \mathbb{R} \tag{III.27}$$

is continuous jointly in the variables $(x^{n-1}, y^{n-1}) \in \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1}$.

Then the following hold.

A1) Let $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and $\big\{\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|x^n) : \alpha = 1, 2, \dots\big\} \subset \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$. Then the joint measure $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{\alpha}_{0,n})(dx^n, dy^n) \xrightarrow{w} (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{0}_{0,n})(dx^n, dy^n)$, where $\overrightarrow{Q}^{0}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$.

A2) Let $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and $\big\{\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|x^n) : \alpha = 1, 2, \dots\big\} \subset \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ and define the family of joint measures $\big\{(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{\alpha}_{0,n})(dx^n, dy^n) : \alpha = 1, 2, \dots\big\}$ having marginals $\{\nu^{\alpha}_{0,n} : \alpha = 1, 2, \dots\}$ on $\mathcal{Y}_{0,n}$ and $\{\mu^{\alpha}_{0,n} : \alpha = 1, 2, \dots\}$ on $\mathcal{X}_{0,n}$. Then $\nu^{\alpha}_{0,n}(dy^n) \xrightarrow{w} \nu^{0}_{0,n}(dy^n)$ and $\mu^{\alpha}_{0,n}(dx^n) \xrightarrow{w} \mu^{0}_{0,n}(dx^n)$, where $\nu^{0}_{0,n} \in \mathcal{M}_1(\mathcal{Y}_{0,n})$ and $\mu^{0}_{0,n} \in \mathcal{M}_1(\mathcal{X}_{0,n})$ are the marginals of $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{0}_{0,n})(dx^n, dy^n)$.

A3) The set of measures $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ is weakly compact.

A4) The set of measures $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ is weakly compact.

A5) Let $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$, $\big\{\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|x^n) : \alpha = 1, 2, \dots\big\} \subset \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$, and $\{\nu^{\alpha}_{0,n} : \alpha = 1, 2, \dots\}$ are the marginals of $\big\{(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{\alpha}_{0,n})(dx^n, dy^n) : \alpha = 1, 2, \dots\big\}$. Then $\overrightarrow{\Pi}^{\alpha}(dx^n, dy^n) \equiv \overleftarrow{P}_{0,n}(dx^n|dy^{n-1}) \otimes \nu^{\alpha}_{0,n}(dy^n) \xrightarrow{w} \overleftarrow{P}_{0,n}(dx^n|dy^{n-1}) \otimes \nu^{0}_{0,n}(dy^n) \equiv \overrightarrow{\Pi}^{0}(dx^n, dy^n)$, where $\nu^{0}_{0,n} \in \mathcal{M}_1(\mathcal{Y}_{0,n})$ is the limit of $\nu^{\alpha}_{0,n} \in \mathcal{M}_1(\mathcal{Y}_{0,n})$.

Part B. *Let $\mathcal{X}_{0,n}$ be a compact Polish space and $\mathcal{Y}_{0,n}$ a Polish space. Assume $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ satisfy the following condition.*

CB*: For all $h(\cdot) \in BC(\mathcal{Y}_{0,n})$, the function*

$$(x^{n}, y^{n-1}) \in \mathcal{X}_{0,n} \times \mathcal{Y}_{0,n-1} \longmapsto \int_{\mathcal{Y}_n} h(y) q_n(dy; y^{n-1}, x^n) \in \mathbb{R} \tag{III.28}$$

is continuous jointly in the variables $(x^n, y^{n-1}) \in \mathcal{X}_{0,n} \times \mathcal{Y}_{0,n-1}$.

Then the following hold.

B1) Let $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ *and* $\{\overleftarrow{P}^{\alpha}_{0,n}(\cdot|y^{n-1}) : \alpha = 1, 2, \dots\} \subset \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$. *Then, the joint measure* $(\overleftarrow{P}^{\alpha}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) \xrightarrow{w} (\overleftarrow{P}^{0}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n)$, *where* $\overleftarrow{P}^{0}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$.

B2) Let $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ *and* $\{\overleftarrow{P}^{\alpha}_{0,n}(\cdot|y^{n-1}) : \alpha = 1, 2, \dots\} \subset \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ *and define the family of joint measures* $\{(\overleftarrow{P}^{\alpha}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) : \alpha = 1, 2, \dots\}$ *having marginals* $\{\nu^{\alpha}_{0,n} : \alpha = 1, 2, \dots\}$ *on* $\mathcal{Y}_{0,n}$ *and* $\{\mu^{\alpha}_{0,n} : \alpha = 1, 2, \dots\}$ *on* $\mathcal{X}_{0,n}$. *Then* $\nu^{\alpha}_{0,n}(dy^n) \xrightarrow{w} \nu^{0}_{0,n}(dy^n)$ *and* $\mu^{\alpha}_{0,n}(dx^n) \xrightarrow{w} \mu^{0}_{0,n}(dx^n)$, *where* $\nu^{0}_{0,n} \in \mathcal{M}_1(\mathcal{Y}_{0,n})$ *and* $\mu^{0}_{0,n} \in \mathcal{M}_1(\mathcal{X}_{0,n})$ *are the marginals of* $(\overleftarrow{P}^{0}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n)$.

B3) The set of measures $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ *is weakly compact.*

B4) The set of measures $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ *is weakly compact.*

B5) Let $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$, $\{\overleftarrow{P}^{\alpha}_{0,n}(\cdot|y^{n-1}) : \alpha = 1, 2, \dots\} \subset \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$, *and* $\{\mu^{\alpha}_{0,n} : \alpha = 1, 2, \dots\}$ *are the marginals of* $\{(\overleftarrow{P}^{\alpha}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) : \alpha = 1, 2, \dots\}$. *Then* $\overleftarrow{\Pi}^{\alpha}(dx^n, dy^n) \equiv \overrightarrow{Q}_{0,n}(dy^n|dx^n) \otimes \mu^{\alpha}_{0,n}(dx^n) \xrightarrow{w} \overrightarrow{Q}_{0,n}(dy^n|dx^n) \otimes \mu^{0}_{0,n}(dx^n) \equiv \overleftarrow{\Pi}^{0}(dx^n, dy^n)$ *where* $\mu^{0}_{0,n} \in \mathcal{M}_1(\mathcal{X}_{0,n})$ *is the limit of* $\mu^{\alpha}_{0,n} \in \mathcal{M}_1(\mathcal{X}_{0,n})$.

Proof. **Part A.** For $\alpha = 1, 2, \dots$, let $\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ be a sequence of forward channels and $(X^{n,(\alpha)}, Y^{n,(\alpha)})$, $n \in \mathbb{N}$ a sequence of the basic joint process corresponding to the backward channel $\overleftarrow{P}_{0,n}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and forward channel $\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$. The important steps for the derivation of A1), A2) are given in [21] for stochastic control problems with randomized controls. Since we shall use A1), A2), and parts of their derivations to show A3)–A5), first we give the details of the derivation of A1) and A2) in Appendix VII-C.

A3) Next, we show that the family of probability measures $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ is weakly compact. By consistency condition $\mathbf{C1}$, any $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ uniquely defines a family of regular conditional distributions $p_i(\cdot; x^{i-1}, y^{i-1}) \in \mathcal{Q}(\mathcal{X}_i; \mathcal{X}_{0,i-1} \times \mathcal{Y}_{0,i-1})$, $i = 0, 1, \dots, n$ via (II.1). Hence, (II.1) can be used to relate weak compactness of the family $p_i(\cdot; x^{i-1}, y^{i-1}) \in \mathcal{Q}(\mathcal{X}_i; \mathcal{X}_{0,i-1} \times \mathcal{Y}_{0,i-1})$, $i = 0, 1, \dots, n$ to weak compactness of the family of measures $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$.

By recalling the derivation in Appendix VII-C, A2), condition (VII.5), for $K_{0,n} = \times_{i=0}^{n} K_i$,

$K_i \in \mathcal{B}(\mathcal{X}_i)$ compact sets, $0 \leq i \leq n$, then

$$\begin{aligned}
\mathbf{P}(K_{0,n}|\mathbf{y}) &\stackrel{\triangle}{=} \int_{K_0} p_0(dx_0) \int_{K_1} p_1(dx_1; x^0, y^0) \dots \int_{K_n} p_n(dx_n; x^{n-1}, y^{n-1}) \\
&\geq \left(1 - \frac{\epsilon_1}{2^{n+1}}\right) \int_{K_0} p_0(dx_0) \int_{K_1} p_1(dx_1; x^0, y^0) \dots \int_{K_{n-1}} p_{n-1}(dx_{n-1}; x^{n-2}, y^{n-2}) \\
&\geq \left(1 - \frac{\epsilon_1}{2^{n+1}}\right)\left(1 - \frac{\epsilon_1}{2^n}\right) \int_{K_0} p_0(dx_0) \int_{K_1} p_1(dx_1; x^0, y^0) \dots \int_{K_{n-2}} p_{n-2}(dx_{n-2}; x^{n-3}, y^{n-3}) \\
&= \left(1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_1}{2^n} + \frac{\epsilon_1^2}{2^{2n+1}}\right) \int_{K_0} p_0(dx_0) \int_{K_1} p_1(dx_1; x^0, y^0) \dots \int_{K_{n-2}} p_{n-2}(dx_{n-2}; x^{n-3}, y^{n-3}) \\
&\geq \left(1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_1}{2^n}\right) \int_{K_0} p_0(dx_0) \int_{K_1} p_1(dx_1; x^0, y^0) \dots \int_{K_{n-2}} p_{n-2}(dx_{n-2}; x^{n-3}, y^{n-3}).
\end{aligned}$$

By repeating the above procedure the following bound is obtained.

$$\begin{aligned}
\mathbf{P}(K_{0,n}|\mathbf{y}) &\geq 1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_1}{2^n} - \frac{\epsilon_1}{2^{n-1}} - \dots - \frac{\epsilon_1}{2^1} \\
&= 1 - \epsilon_1 \sum_{i=1}^{n} \frac{1}{2^{i+1}} \\
&\geq 1 - \epsilon_1, \qquad \text{for any } n \in \mathbb{N} \text{ and for every } \overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1}).
\end{aligned}$$

Since $\{K_i : i = 0, 1, \dots, n\}$ are compact, from the last inequality it follows that the family of measures $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ it tight, and hence weakly compact. This completes the derivation of A3).

A4) Weak compactness of the family of measures $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ follows from the fact that $\mathcal{Y}_{0,n}$ is a compact Polish space. Indeed, any $\overrightarrow{Q}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ uniquely defines a family of regular conditional distributions $q_i(\cdot; y^{i-1}, x^i) \in \mathcal{Q}(\mathcal{Y}_i; \mathcal{Y}_{0,i-1} \times \mathcal{X}_{0,i})$, $i = 0, 1, \dots, n$. It was already shown that given $\epsilon_2 > 0$ one can construct compact sets $\Phi_0 \subset \mathcal{Y}_0, \Phi_1 \subset \mathcal{Y}_1, \dots, \Phi_{n-1} \subset \mathcal{Y}_{n-1}$, such that (see Appendix VII-C, A1)) $q_n(\Phi_n; y^{n-1}, x^n) \geq 1 - \frac{\epsilon_2}{2^{n+1}}$, for any $y_0 \in \Phi_0, y_1 \in \Phi_1, \dots, y_{n-1} \in \Phi_{n-1}, x^n \in \mathcal{X}_{0,n}$. Utilizing the last expression and the unique representation of $\overrightarrow{Q}_{0,n}(\cdot|x^n)$ given by (II.4), tightness of $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ can be shown, and hence it is weakly compact. This completes the derivation of A4).

A5) Utilizing the weak convergence $\nu^{\alpha}_{0,n} \xrightarrow{w} \nu^{0}_{0,n}$ (shown under Appendix VII-C, A2)), we shall show weak convergence of the convolution of measures $\overrightarrow{\Pi}^{\alpha}_{0,n}(dx^n, dy^n) \equiv \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \otimes \nu^{\alpha}_{0,n}(dy^n) \xrightarrow{w} \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \otimes \nu^{0}_{0,n}(dy^n) \equiv \overrightarrow{\Pi}^{0}_{0,n}(dx^n, dy^n)$, when $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ is fixed. We show weak convergence by considering integrals with respect to $g_{0,n}(x^n)h_{0,n}(y^n)$,

where $g_{0,n}(\cdot) \in BC(\mathcal{X}_{0,n})$ and $h_{0,n}(\cdot) \in BC(\mathcal{Y}_{0,n})$. Let $\epsilon > 0$ be given. Condition **CA** implies that the function mapping

$$y^{n-1} \in \mathcal{Y}_{0,n-1} \longmapsto \int_{\mathcal{X}_{0,n}} g(x^n) \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \in \mathbb{R} \tag{III.29}$$

is continuous. Hence, by the weak convergence $\nu^{\alpha}_{0,n} \xrightarrow{w} \nu^{0}_{0,n}$ and the continuity of the function mapping (III.29) then there exists $N \in \mathbb{N}$ such that for all $\alpha \geq N$

$$\left| \int_{\mathcal{Y}_{0,n}} \left(\int_{\mathcal{X}_{0,n}} g(x^n) \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \right) h(y^n) \nu^{0}_{0,n}(dy^n) \right.$$
$$\left. - \int_{\mathcal{Y}_{0,n}} \left(\int_{\mathcal{X}_{0,n}} g(x^n) \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \right) h(y^n) \nu^{\alpha}_{0,n}(dy^n) \right| \leq \epsilon.$$

Since $\epsilon > 0$ is arbitrary, then the derivation of A5) is complete.

Part B. The methodology is similar to that of **Part A.**, hence it is omitted. □

It is important to illustrate how some parts of Theorem III.5 compare to analogous results for mutual information discussed in [20], [27]. To this end, consider **Part B.**, B1). If we use mutual information [20, Lemma 2], then the sequence of joint measures is defined by $P^{\alpha}_{X^n,Y^n}(dx^n, dy^n) \stackrel{\triangle}{=} P_{Y^n|X^n}(dy^n|x^n) \otimes P^{\alpha}_{X^n}(dx^n)$, and this involves a single convolution, since $P_{X^n}(dx^n)$ is not conditioned on $y^n \in \mathcal{Y}_{0,n}$. Clearly, if the mapping $x^n \longrightarrow P_{Y^n|X^n}(\cdot|x^n)$ is weakly continuous, and $P^{\alpha}_{X^n}(dx^n)$ converges weakly to $P^{0}_{X^n}(x^n)$, then $P^{\alpha}_{X^n,Y^n}(dx^n, dy^n)$ converges weakly to $P_{Y^n|X^n}(dy^n|x^n) \otimes P^{0}_{X^n}(dx^n) = P^{0}_{X^n,Y^n}(dx^n, dy^n)$, and so does its marginal on $\mathcal{Y}_{0,n}$. On the other hand, if we use directed information, then the joint measure $P_{X^n,Y^n}(dx^n, dy^n) \stackrel{\triangle}{=} \otimes_{i=0}^{n} P_{Y_i|Y^{i-1},X^i}(dy_i|y^{i-1}, x^i) \otimes P_{X_i|X^{i-1},Y^{i-1}}(dx_i|x^{i-1}, y^{i-1})$ involves an $(n+1)$-fold convolution, and $P_{X_i|X^{i-1},Y^{i-1}}(\cdot|\cdot,\cdot)$ is a function of $y^{n-1} \in \mathcal{Y}_{0,n-1}$, hence a significant level of additional complexity incurs compared to mutual information. Nevertheless, condition **CB** is the natural generalization of the weak continuity of the mapping $x^n \longrightarrow P_{Y^n|X^n}(\cdot|x^n)$ assumed for the mutual information by Csiszár in [20], to causally conditioned $(n+1)$-fold convolutional measures.

Theorem III.5 is important for several extremum problems involving directed information. In the next Remark we discuss possible applications of the Theorem III.5.

Remark III.6.

1) Consider extremum problems of capacity of channels with memory and feedback. Then, in the absence of input constraints, the aim is to find the input conditional distribution $\overleftarrow{P}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$*, which achieves the infimum of directed information. To show that such a conditional distribution exists, it is sufficient to show compactness of* $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$*, and A3) establishes weak compactness. Moreover, in the presence of power constraints* $\overleftarrow{P}(\cdot|y^{n-1}) \in \mathcal{P}_{0,n}(P) \subset \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$*, to show that such conditional distributions exist, by Prohorov's theorem (Appendix VI-A, Theorem VI.5), it is sufficient to show* $\mathcal{P}_{0,n}(P)$ *is closed and uniformly tight subset of the weakly compact set* $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$*.*

2) Consider extremum problems of sequential or nonanticipative lossy data compression with distortion constraint $\mathcal{Q}_{0,n}(D) \subset \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ *[4], [33]. Then Theorem III.5,* **Part A.**, *A4) is important to show weak compactness of* $\mathcal{Q}_{0,n}(D)$ *as a subset of the weakly compact set* $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$*, utilizing Prohorov's theorem (Appendix VI-A, Theorem VI.4). For such problems, condition* **CA** *is replaced by*

$$x^{n-1} \in \mathcal{X}_{0,n-1} \longmapsto \int_{\mathcal{X}_n} g(x) p_n(dx; x^{n-1}) \in \mathbb{R}$$

is continuous in $x^{n-1} \in \mathcal{X}_{0,n-1}$*, since the source* $\{p_n(x_n; x^{n-1}) : \ n \in \mathbb{N}\}$ *is not affected by past reconstructions* $\{y_n : n \in \mathbb{N}\}$ *(see [4]).*

3) Consider extremum problems of capacity for a class of channels with memory and feedback, such as, arbitrary varying channels [20]. Such problems are defined by the max-min operations of directed information, where the minimizer is over the class of channels [34]. To investigate such capacity problems one has to establish coding theorems, and showing compactness over the class of channel conditional distributions, in addition to channel input distributions is critical. Theorem III.5, **Part B.**, *B3) establishes weak compactness of this family of channels.*

4) Consider extremum problems of sequential or nonanticipative lossy data compression for a class of sources. Then such problems are defined by min-max operations of directed information, where the maximizer is over the class of source distributions [35]. To investigate such data compression problems one has to establish coding theorems, and to show compactness over the class of source distributions, in addition to the reproduction distributions, Theorem III.5, **Part A.**, *A3) is crucial.*

In the previous applications, Theorem III.5 gives the choice of considering either $\mathcal{Y}_{0,n}$ *or* $\mathcal{X}_{0,n}$ *to be compact.*

D. Lower Semicontinuity of Directed Information

We are now ready to utilize the results of Theorem III.5, to show lower semicontinuity of directed information $I(X^n \to Y^n) \equiv \mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$. This may be viewed as a generalization of lower semicontinuity of mutual information $I(X^n;Y^n) \equiv \mathbb{I}_{X^n;Y^n}(P_{X^n}, Q_{Y^n|X^n})$, with respect to P_{X^n} for fixed $Q_{Y^n|X^n}$, and with respect to $Q_{Y^n|X^n}$ for fixed P_{X^n}.

Theorem III.7.

1) Suppose the conditions in Theorem III.5, **Part A.**, *hold.*
Then $\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ *is lower semicontinuous on* $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ *for fixed* $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$.
2) Suppose the conditions in Theorem III.5, **Part B.**, *hold.*
Then $\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ *is lower semicontinuous on* $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ *for fixed* $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$.

Proof. 1) We need to show that for any sequence $\{\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|x^n) : \alpha = 1, 2, \ldots\}$, $\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ such that $\overrightarrow{Q}^{\alpha}_{0,n}(\cdot|x^n) \xrightarrow{w} \overrightarrow{Q}^{0}_{0,n}(\cdot|x^n)$ then

$$\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}^{0}_{0,n}) \leq \liminf_{\alpha \to \infty} \mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}^{\alpha}_{0,n}).$$

Define the sequence $P^{\alpha}_{0,n}(dx^n, dy^n) \stackrel{\triangle}{=} (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{\alpha}_{0,n})(dx^n, dy^n)$. Weak convergence $P^{\alpha}_{0,n}(dx^n, dy^n) \xrightarrow{w} (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{0}_{0,n})(dx^n, dy^n) \equiv P^{0}_{0,n}(dx^n, dy^n)$ is shown by considering integrals with respect to a test function $\phi_{0,n}(\cdot,\cdot) \in BC(\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n})$ via

$$\int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \phi_{0,n}(x^n, y^n) P^{\alpha}_{0,n}(dx^n, dy^n) = \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \phi_{0,n}(x^n, y^n) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{\alpha}_{0,n})(dx^n, dy^n).$$

By Theorem III.5, **Part A.**, A1), $P^{\alpha}_{0,n}(dx^n, dy^n) \xrightarrow{w} P^{0}_{0,n}(dx^n, dy^n)$. Similarly, $\overrightarrow{\Pi}^{\alpha}_{0,n} \stackrel{\triangle}{=} \overleftarrow{P}_{0,n} \otimes \nu^{\alpha}_{0,n}$, where $\{\nu^{\alpha}_{0,n} : \alpha = 1, 2, \ldots\}$ are the marginals of $\{P^{\alpha}_{0,n} : \alpha = 1, 2, \ldots\}$, and by Theorem III.5, **Part A.**, A5) we have

$$\overrightarrow{\Pi}^{\alpha}_{0,n} = \overleftarrow{P}_{0,n} \otimes \nu^{\alpha}_{0,n} \xrightarrow{w} \overrightarrow{\Pi}^{0}_{0,n} = \overleftarrow{P}_{0,n} \otimes \nu^{0}_{0,n}.$$

Recall the definition of directed information via relative entropy given by

$$\mathbb{D}(P_{0,n} || \overrightarrow{\Pi}_{0,n}) = \mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n} || \overleftarrow{P}_{0,n} \otimes \nu_{0,n}) = \mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}). \tag{III.30}$$

Relative entropy is lower semicontinuous, hence

$$\mathbb{D}(P^0_{0,n}||\overrightarrow{\Pi}^0_{0,n}) = \mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^0_{0,n}||\overrightarrow{\Pi}^0_{0,n}) \leq \liminf_{\alpha\to\infty} \mathbb{D}(P^\alpha_{0,n}||\overrightarrow{\Pi}^\alpha_{0,n}). \tag{III.31}$$

By (III.30) it follows that (III.31) is also equivalent to

$$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}^0_{0,n}) \leq \liminf_{\alpha\to\infty} \mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}^\alpha_{0,n})$$

Hence, directed information is lower semicontinuous as a functional of $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ for a fixed $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. This completes the derivation of 1).

2) The derivation is similar to 1). □

By comparison with lower semicontinuity of mutual information $I(X^n;Y^n) \equiv \mathbb{I}_{X^n;Y^n}(P_{X^n}, Q_{Y^n|X^n})$, it is clear that directed information imposes additional assumptions for its derivation (e.g., those given in Theorem III.5).

E. Continuity of Directed Information

Note that Theorem III.5 together with Theorem III.7 are important to establish existence of the optimal reconstruction distribution for sequential and nonanticipative rate distortion functions (by utilizing Weierstrass' Theorem). However, many problems in information theory involve extremum problems defined as maximizations of directed information, with respect to the feedback channels $\{p_i(dx_i; x^{i-1}, y^{i-1}) : \ i = 0, 1, \ldots, n\}$, and for such problems it is desirable to have upper semicontinuity of directed information with respect to $\overleftarrow{P}_{0,n}$. Since by Theorem III.7, directed information is lower semicontinuous with respect to $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$, to investigate extremum problems involving feedback capacity, it is sufficient to show continuity of the functional $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ with respect to $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ for a fixed $\overrightarrow{Q}_{0,n}$. Continuity of mutual information based on single letter expression is shown in [20, Lemma 7], and in [36, Theorem 3.2] under weaker conditions. Here, we show continuity of directed information by following the procedure in [36], generalized to the directed information functional $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$. First, we shall need the following Lemma.

Lemma III.8.

For a given $\overleftarrow{P}_{0,n} \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ *and* $\overrightarrow{Q}_{0,n} \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ *define*

$$\Big|\mathbb{I}_{X^n\to Y^n}\Big|(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}) \triangleq \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \left|\log\left(\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}\right)\right| d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}).$$

Then the following inequalities hold.

$$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})\leq|\mathbb{I}_{X^n\to Y^n}|(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})\leq\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})+\frac{2}{e\ln 2}. \tag{III.32}$$

Proof. The directed information as it is defined in Theorem III.1 and (III.14) can be expressed as follows.

$$\begin{aligned}\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n}) &\stackrel{\triangle}{=}\mathbb{D}(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}||\overrightarrow{\Pi}_{0,n})\\ &=\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\left(\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}\right)d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})\\ &=\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\left(\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}\right)\left(\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}\right)d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n}).\end{aligned} \tag{III.33}$$

The first inequality in (III.32) is obvious. For the second inequality recall the inequality [29, Section 2.3, p. 13] $-\frac{1}{e\ln 2}\leq x\log_2 x$, $x\in[0,\infty)$, $0\log 0$ is assumed to be 0. Then,

$$|x\log_2 x|\leq x\log_2 x+\frac{2}{e\ln 2}. \tag{III.34}$$

Using the last inequality in (III.33) with $x=\left(\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{P}_{0,n}\otimes\nu_{0,n})}\right)$, establishes the second inequality in (III.32). □

Now, we are ready to state the Theorem which establishes continuity with respect to weak convergence of $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})$ for a fixed $\overrightarrow{Q}_{0,n}$, as a functional of $\overleftarrow{P}_{0,n}\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$.

Theorem III.9.

Consider a forward channel $\overrightarrow{Q}_{0,n}(\cdot|x^n)\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$, and a closed family of feedback channels $\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\subseteq\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. Suppose there exists a family of measures $\bar{\nu}_{0,n}(dy^n)$ on $(\mathcal{Y}_{0,n},\mathcal{B}(\mathcal{Y}_{0,n}))$ such that $\overrightarrow{Q}_{0,n}(\cdot|x^n)\ll\bar{\nu}_{0,n}(dy^n)$ with RNDs $\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\stackrel{\triangle}{=}\frac{\overrightarrow{Q}_{0,n}(\cdot|x^n)}{\bar{\nu}_{0,n}(\cdot)}(y^n)$. Furthermore, suppose the following conditions hold.

A) The family of RNDs $\xi_{\bar{\nu}_{0,n}}(x^n,y^n)$ is continuous on $\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}$, and $\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)$ is uniformly integrable over $\left\{\bar{\nu}_{0,n}\otimes\overleftarrow{P}_{0,n}:\overleftarrow{P}_{0,n}\in\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\right\}$.

B) For a fixed $y^n\in\mathcal{Y}_{0,n}$, the RND $\xi_{\bar{\nu}_{0,n}}(x^n,y^n)$ is uniformly integrable over $\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. Then, $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})$ as a functional of $\overleftarrow{P}_{0,n}\in\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$, $\overrightarrow{Q}_{0,n}\}\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ is bounded and weakly continuous over $\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$.

Proof. The derivation is shown in Appendix VII-D. □

Note that Theorem III.5 gives conditions for $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ to be weakly compact, hence under these conditions, and the additional conditions A), B), of Theorem III.9, then $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})$ is bounded weakly continuous over $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. Thus, Theorem III.5 together with Theorem III.9 establish all prerequisites to address extremum problems of capacity of channels with memory and feedback described under Remark III.6. Two such problems are briefly discussed in [37].

F. Extension of Directed Information to Arbitrary sequences of RV's

In this section, we demonstrate how the previous results are easily generalized to three, or more, sequences of RV's. These extensions have implications in networks communication, and in communications with side information at either the transmitter or the receiver [25], [26].
To facilitate the demonstration, we consider the following case first.

Case 1: The sequence of RV's $X^n\in\mathcal{X}_{0,n}$ is defined by $X^n=(X^{1,n},X^{2,n})\in\mathcal{X}^1_{0,n}\times\mathcal{X}^2_{0,n}\equiv\mathcal{X}_{0,n}$, where $X^{1,n}=\{X^1_i:\ i=0,1,\ldots,n\}$ and $X^{2,n}=\{X^2_i:\ i=0,1,\ldots,n\}$.

Then the two sequences of conditional distributions are $\{p^1_i(dx^1_i,dx^2_i;x^{1,i-1},x^{2,i-1},y^{i-1}):i=0,1,\ldots,n\}$ and $\{q^1_i(dy^1_i;y^{1,i-1},x^{1,i},x^{2,i}):i=0,1,\ldots,n\}$, respectively. Consequently, all construction of consistent families of conditional distributions and the results obtained so far extend naturally to directed information $\mathbb{I}_{(X^{1,n},X^{2,n})\to Y^n}(\overleftarrow{P}^1_{0,n},\overrightarrow{Q}^1_{0,n})$, where $\overleftarrow{P}^1_{0,n}(dx^{1,n},dx^{2,n}|y^{n-1})=\otimes_{i=0}^n p^1_i(dx^1_i,dx^2_i;x^{1,i-1},x^{2,i-i},y^{i-1})$, and $\overrightarrow{Q}^1_{0,n}(dy^n|x^{1,n},x^{2,n})=\otimes_{i=0}^n q^1_i(dy_i;y^{i-1},x^{1,i},x^{2,i})$.
Next, we consider the following case.

Case 2: The sequence of RV's $Y^n\in\mathcal{Y}_{0,n}$ is defined by $Y^n=(Y^{1,n},Y^{2,n})\in\mathcal{Y}^1_{0,n}\times\mathcal{Y}^2_{0,n}\equiv\mathcal{Y}_{0,n}$, where $Y^{1,n}=\{Y^1_i:\ i=0,1,\ldots,n\}$ and $Y^{2,n}=\{Y^2_i:\ i=0,1,\ldots,n\}$.

Then the two sequences of conditional distributions are $\{p^2_i(dx_i;x^{i-1},y^{1,i-1},y^{2,i-1}):i=0,1,\ldots,n\}$ and $\{q^2_i(dy^1_i,dy^2_i;y^{1,i-1},y^{2,i-1},x^i):i=0,1,\ldots,n\}$, respectively. Consequently, all construction of consistent families of conditional distributions and the results obtained so far extend naturally to directed information $\mathbb{I}_{X^n\to(Y^{1,n},Y^{2,n})}(\overleftarrow{P}^2_{0,n},\overrightarrow{Q}^2_{0,n})$, where $\overleftarrow{P}^2_{0,n}(dx^n|y^{1,n-1},y^{2,n-1})=\otimes_{i=0}^n p^1_i(dx_i;x^{i-1},y^{1,i-i},y^{2,i-1})$, and $\overrightarrow{Q}^1_{0,n}(dy^{1,n},dy^{2,n}|x^n)=\otimes_{i=0}^n q^1_i(dy^1_i,dy^2_i;y^{1,i-1},y^{2,i-1},x^i)$.
Clearly, **Case 1** and **Case 2** can be generalized to an arbitrary number of sequences of RV's.

IV. Variational Equalities of Directed Information

In this section we generalize the well-known variational equalities of mutual information [19] to directed information. For single letter mutual information these are discussed in [38]. Before we pursue this task, we recall the variational equalities of mutual information. It is well-known that mutual information $I(X^n;Y^n) \equiv \mathbb{I}_{X^n;Y^n}(P_{X^n}, P_{Y^n|X^n})$ can be expressed as the minimization or maximization of relative entropy functionals as follows [19].

Min: Given a reconstruction distribution $P_{Y^n|X^n}(dy^n|x^n)$, a source P_{X^n}, and any distribution $V_{Y^n}(dy^n)$ then

$$\mathbb{I}_{X^n;Y^n}(P_{X^n}, P_{Y^n|X^n}) = \inf_{V_{Y^n} \in \mathcal{M}_1(\mathcal{Y}_{0,n})} \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log \left(\frac{P_{Y^n|X^n}(dy^n|x^n)}{V_{Y^n}(dy^n)} \right) P_{Y^n|X^n}(dy^n|x^n) P_{X^n}(dx^n) \tag{IV.1}$$

where the infimum is achieved at $V_{Y^n}(dy^n) = \int_{\mathcal{X}_{0,n}} P_{Y^n|X^n}(dy^n|x^n) \mu_{X^n}(dx^n)$.

Max: Given a channel $P_{Y^n|X^n}(dy^n|x^n)$, a distribution $P_{X^n}(dx^n)$, and any conditional distribution $P_{X^n|Y^n}(dx^n|y^n)$ then

$$\mathbb{I}_{X^n;Y^n}(P_{X^n}, P_{Y^n|X^n}) = \sup_{\substack{P_{X^n|Y^n} \in \\ \mathcal{Q}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n})}} \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log \left(\frac{P_{X^n|Y^n}(dx^n|y^n)}{P_{X^n}(dx^n)} \right) P_{Y^n|X^n}(dy^n|x^n) P_{X^n}(dx^n) \tag{IV.2}$$

and the supremum is achieved at $P_{X^n|Y^n}(dx^n|y^n) = \frac{P_{Y^n|X^n}(dy^n|x^n) \otimes P_{X^n}(dx^n)}{\int_{\mathcal{X}_{0,n}} P_{Y^n|X^n}(dy^n|x^n) \otimes P_{X^n}(dx^n)}$.

Both variational equalities are used in the Blahut-Arimoto algorithm [19], [39] to derive an iterative computational scheme for capacity and rate distortion via max-min and min-max operations, respectively. We shall derive analogous variational equalities for directed information. Before we state the main theorem, we introduce some preliminaries.

Let $\mathbf{P}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$, and let $P_{0,n}(dx^n, dy^n) = \overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \otimes \overrightarrow{Q}_{0,n}(dy^n|x^n)$ be the given distribution constructed from the basic feedback channel $\mathbf{P}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ and forward channel $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$ (by projection onto finite number of coordinates).

Let $\mathbf{S}(\cdot|\mathbf{x})$ be any measure on $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ depending parametrically on $\mathbf{x} \in \mathcal{X}^{\mathbb{N}}$ satisfying the

consistency condition.

C3: If $F \in \mathcal{B}(\mathcal{Y}_{0,n})$, then $\mathbf{S}(F|\mathbf{x})$ is a $\mathcal{B}(\mathcal{X}_{0,n-1})-$measurable.

Denote this family of measures by $\mathbf{S}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C3}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$. By Remark II.1, for any family of measures $\mathbf{S}(\cdot|\mathbf{x})$ on $(\mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ satisfying consistency condition **C3**, there exists a collection of stochastic kernels $\{s_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n-1}) : n \in \mathbb{N}\}$ connected to $\mathbf{S}(\cdot|\mathbf{x})$. Indeed, given a cylinder set $D \in \mathcal{B}(\mathcal{Y}_{0,n})$ of the form

$$D \stackrel{\triangle}{=} \{\mathbf{y} \in \mathcal{Y}^{\mathbb{N}} : y_0 \in D_0, y_1 \in D_1, \ldots, y_n \in D_n\},\ D_i \in \mathcal{B}(\mathcal{Y}_i),\ 0 \leq i \leq n$$

then

$$\begin{aligned}\mathbf{S}(D|\mathbf{x}) &= \int_{D_0} s_0(dy_0) \int_{D_1} s_1(dy_1; y_0, x_0) \ldots \int_{D_n} s_n(dy_n; y^{n-1}, x^{n-1}) \\ &\equiv \overleftarrow{S}_{0,n}(\times_{i=0}^{n} D_i | x^{n-1}). \end{aligned} \tag{IV.3}$$

Unlike the measure $\overrightarrow{Q}_{0,n}(\cdot|x^n)$ which is conditioned on $x^n \in \mathcal{X}_{0,n}$, the measure $\overleftarrow{S}_{0,n}(\cdot|x^{n-1})$ is conditioned on $x^{n-1} \in \mathcal{X}_{0,n-1}$.

Let $\mathbf{R}(\cdot|\mathbf{y})$ be any family of measures on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ depending parametrically on $\mathbf{y} \in \mathcal{Y}^{\mathbb{N}}$ satisfying the consistency condition.

C4: If $E \in \mathcal{B}(\mathcal{X}_{0,n})$, then $\mathbf{R}(E|\mathbf{y})$ is a $\mathcal{B}(\mathcal{Y}_{0,n})-$measurable.

Denote this family of measures by $\mathbf{R}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C4}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$. Similarly as before, by Remark II.1, for any family of measures $\mathbf{R}(\cdot|\mathbf{y})$ on $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ satisfying consistency condition **C4**, there exists a collection of stochastic kernels $\{r_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n}) : n \in \mathbb{N}\}$ connected to $\mathbf{R}(\cdot|\mathbf{y})$. Thus, given a cylinder set $E \in \mathcal{B}(\mathcal{X}_{0,n})$ of the form

$$E \stackrel{\triangle}{=} \{\mathbf{x} \in \mathcal{X}^{\mathbb{N}} : x_0 \in E_0, x_1 \in E_1, \ldots, x_n \in E_n\},\ E_i \in \mathcal{B}(\mathcal{X}_i),\ 0 \leq i \leq n$$

then

$$\begin{aligned}\mathbf{R}(E|\mathbf{y}) &= \int_{E_0} r_0(dx_0; y_0) \int_{E_1} r_1(dx_1; x_0, y^1) \ldots \int_{E_n} r_n(dx_n; x^{n-1}, y^n) \\ &\equiv \overrightarrow{R}_{0,n}(\times_{i=0}^{n} E_i | y^n). \end{aligned} \tag{IV.4}$$

Note the difference between the stochastic kernels $\{p_i(dx_i; x^{i-1}, y^{i-1}) : i \in \mathbb{N}\}$ which define $\overleftarrow{P}_{0,n}(dx^n|y^{n-1})$, and the stochastic kernels $\{r_i(dx_i; x^{i-1}, y^i) : \ i \in \mathbb{N}\}$ which define

$\overrightarrow{R}_{0,n}(dx^n|y^n)$. Define their joint distribution $(\overleftarrow{S}_{0,n}\otimes\overrightarrow{R}_{0,n})(dx^n,dy^n)$ on $\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}},\odot_{n\in\mathbb{N}}\mathcal{B}(\mathcal{X}_n)\odot\mathcal{B}(\mathcal{Y}_n)\big)$.

The following theorem gives two variational equalities for directed information, which are analogous to (IV.1), (IV.2).

Theorem IV.1.

Let $\mathbf{P}(\cdot|\mathbf{y})\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ *and* $\mathbf{Q}(\cdot|\mathbf{x})\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$*, and construct from them the measures* $P_{0,n}(dx^n,dy^n)=(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n)$, $\overrightarrow{\Pi}(dx^n,dy^n)=\overleftarrow{P}_{0,n}(dx^n|y^{n-1})\otimes\nu_{0,n}(dy^n)$, $\nu_{0,n}(dy^n)=P_{0,n}(\mathcal{X}_{0,n},dy^n)$*. Then we have the following variational equalities.*

Part A. *For any arbitrary measure* $\bar{\nu}_{0,n}\in\mathcal{M}_1(\mathcal{Y}_{0,n})$ *we have*

$$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})\stackrel{\triangle}{=}\mathbb{D}(P_{0,n}||\overrightarrow{\Pi}_{0,n})$$

$$=\inf_{\bar{\nu}_{0,n}\in\mathcal{M}_1(\mathcal{Y}_{0,n})}\mathbb{D}(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}||\overleftarrow{P}_{0,n}\otimes\bar{\nu}_{0,n}) \tag{IV.5}$$

$$=\inf_{\bar{\nu}_{0,n}\in\mathcal{M}_1(\mathcal{Y}_{0,n})}\Big\{\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\bar{\nu}_{0,n}(dy^n)}\Big)(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n)\Big\} \tag{IV.6}$$

and the infimum in (IV.5) is achieved by

$$\bar{\nu}^*_{0,n}(dy^n)=\bar{\nu}_{0,n}(dy^n)=\int_{\mathcal{X}_{0,n}}(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n).$$

Part B. *For any* $\mathbf{S}(\cdot|\mathbf{x})\in\mathcal{Q}^{\mathbf{C3}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$ *and* $\mathbf{R}(\cdot|\mathbf{y})\in\mathcal{Q}^{\mathbf{C4}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ *then*

$$\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})=\mathbb{D}(P_{0,n}||\overrightarrow{\Pi}_{0,n})$$

$$=\sup_{\substack{\overleftarrow{S}_{0,n}\otimes\overrightarrow{R}_{0,n}\in\mathcal{M}_1(\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}):\\ \overleftarrow{S}_{0,n}\in\mathcal{Q}^{\mathbf{C3}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}}),\overrightarrow{R}_{0,n}\in\mathcal{Q}^{\mathbf{C4}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})}}\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\Big(\frac{d(\overleftarrow{S}_{0,n}\otimes\overrightarrow{R}_{0,n})}{d(\overrightarrow{\Pi}_{0,n})}\Big)d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}) \tag{IV.7}$$

where the supremum in (IV.7) is achieved when the RND satisfies

$$\Lambda_{0,n}(x^n,y^n)\stackrel{\triangle}{=}\frac{d(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})}{d(\overleftarrow{S}_{0,n}\otimes\overrightarrow{R}_{0,n})}=1-a.s.,\ n\in\mathbb{N}. \tag{IV.8}$$

Equivalently,

$$\lambda_i(x^i,y^i)\stackrel{\triangle}{=}\frac{p_i(dx_i;x^{i-1},y^{i-1})\otimes q_i(dy_i;y^{i-1},x^i)}{s_i(dy_i;y^{i-1},x^{i-1})\otimes r_i(dx_i;x^{i-1},y^i)}=1-a.s.,\ i=0,1,\ldots,n. \tag{IV.9}$$

Moreover, if $q_i(\cdot; y^{i-1}, x^i) \ll s_i(\cdot; y^{i-1}, x^{i-1})$ *and* $p_i(\cdot; x^{i-1}, y^{i-1}) \ll r_i(\cdot; x^{i-1}, y^i),\ i = 0, 1, \ldots, n$, *then*

$$\otimes_{i=0}^{n} \frac{q_i(dy_i; y^{i-1}, x^i)}{s_i(dy_i; y^{i-1}, x^{i-1})} = \otimes_{i=0}^{n} \left(\frac{p_i(dx_i; x^{i-1}, y^{i-1})}{r_i(dx_i; x^{i-1}, y^i)} \right)^{-1} - a.s.,\ \ n \in \mathbb{N} \tag{IV.10}$$

or equivalently,

$$\frac{q_i(dy_i; y^{i-1}, x^i)}{s_i(dy_i; y^{i-1}, x^{i-1})} = \left(\frac{p_i(dx_i; x^{i-1}, y^{i-1})}{r_i(dx_i; x^{i-1}, y^i)} \right)^{-1} - a.s.,\ \ i = 0, 1, \ldots, n. \tag{IV.11}$$

Proof. **Part A.** From Theorem III.1:

$$\mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n} || \overleftarrow{P}_{0,n} \otimes \bar{\nu}_{0,n}) = \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log \Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\bar{\nu}_{0,n}(dy^n)} \Big) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) \tag{IV.12}$$

$$= \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log \Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)} \Big) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) + \mathbb{D}(\nu_{0,n} || \bar{\nu}_{0,n}) \tag{IV.13}$$

$$\geq \mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n} || \overleftarrow{P}_{0,n} \otimes \nu_{0,n}). \tag{IV.14}$$

Moreover, equality holds in (IV.14) when $\nu_{0,n} = \bar{\nu}_{0,n}$. Hence, $\mathbb{D}(P_{0,n} || \overrightarrow{\Pi}_{0,n})$ in (III.19) can be expressed via variational equalities (IV.5) and (IV.6).

Part B. Consider the difference between $I(X^n \to Y^n) = \mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n} || \overrightarrow{\Pi}_{0,n})$ given by (III.19) and the right hand side of (IV.7) (without the supremum). Then

$$\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}) - \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log \Big(\frac{d(\overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n})}{d(\overleftarrow{P}_{0,n} \otimes \nu_{0,n})} \Big) d(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})$$

$$= \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log \Big(\frac{d(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})}{d(\overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n})} \Big) d(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})$$

$$\geq \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \Big(1 - \frac{\overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n}}{\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n}} \Big) (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) = 0 \tag{IV.15}$$

where (IV.15) follows from the inequality $\log x \geq 1 - \frac{1}{x}$, $x > 0$, which holds with equality if and only if $x = 1$. Furthermore, equality holds in (IV.15) when the RND $\Lambda_{0,n}(x^n, y^n) \stackrel{\triangle}{=} \frac{\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n}}{\overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n}} = 1$, $\overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n} - a.s.$ in (x^n, y^n). Since $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}) = (\overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n})(\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}) = 1$, this condition is equivalent to $\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n} = \overleftarrow{S}_{0,n} \otimes \overrightarrow{R}_{0,n}$. By conditioning (IV.8) on $\mathcal{B}(\mathcal{X}_{0,n-1}) \odot \mathcal{B}(\mathcal{Y}_{0,n-1})$ one obtains (IV.9). Furthermore, (IV.10) is obtained from (IV.8), while (IV.11) is obtained by conditioning. □

Next, we discuss the relation between the variational equalities of directed information given by (IV.7) and the variational equalities of mutual information given by (IV.2). Clearly, (IV.2) is also equivalent to

$$\sup_{P_{X^n|Y^n}\otimes P_{Y^n}} \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\left(\frac{P_{X^n|Y^n}(dx^n|y^n)\otimes P_{Y^n}(dy^n)}{P_{X^n}(dx^n)\times P_{Y^n}(dy^n)}\right) P_{Y^n|X^n}(dy^n|x^n)\otimes P_{X^n}(dx^n) \quad \text{(IV.16)}$$

since the RND in (IV.16) is another version of the one in (IV.2). Thus, (IV.7) is understood in the context of (IV.16), in which the directed information function is utilized together with the decomposition $\overleftarrow{S}_{0,n}\otimes\overrightarrow{R}_{0,n}$. Consider obtaining the analog of the maximizing measure in (IV.2). Suppose $q_i(\cdot;y^{i-1},x^i)\ll s_i(\cdot;y^{i-1},x^{i-1})$, $i=0,1,\ldots,n$. Then from (IV.9) we obtain

$$\begin{aligned} r_i(dx_i;x^{i-1},y^i) &= \left(\frac{q_i(dy_i;y^{i-1},x^i)}{s_i(dy_i;y^{i-1},x^{i-1})}\right)\otimes p_i(dx_i;x^{i-1},y^{i-1}), \qquad i=0,1,\ldots,n \\ &= \frac{q_i(dy_i;y^{i-1},x^i)\otimes p_i(dx_i;x^{i-1},y^{i-1})}{\int_{\mathcal{X}_i} q_i(dy_i;y^{i-1},x^i)\otimes p_i(dx_i;x^{i-1},y^{i-1})}, \qquad i=0,1,\ldots,n. \end{aligned}$$

Consequently,

$$\overrightarrow{R}_{0,n}(dx^n|y^n)=\otimes_{i=0}^n \frac{q_i(dy_i;y^{i-1},x^i)\otimes p_i(dx_i;x^{i-1},y^{i-1})}{\int_{\mathcal{X}_i} q_i(dy_i;y^{i-1},x^i)\otimes p_i(dx_i;x^{i-1},y^{i-1})}, \qquad n\in\mathbb{N}.$$

The previous expression is the analog of the maximizing distribution $P_{X^n|Y^n}$ in (IV.2). Finally, we note that the optimization in (IV.7) can be done by keeping $\overleftarrow{S}_{0,n}$ fixed, generated by $\mathbf{P}(\cdot|\cdot)\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\cdot)\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$.

One can obtain the variational equalities (IV.5), (IV.6) of Theorem IV.1 by using the equivalent definition consisting of a family of functions $\Big\{p_n(\cdot;\cdot,\cdot)\in\mathcal{Q}(\mathcal{X}_n;\mathcal{X}_{0,n-1}\times\mathcal{Y}_{0,n-1}):n\in\mathbb{N}\Big\}$ and $\Big\{q_n(\cdot;\cdot,\cdot)\in\mathcal{Q}(\mathcal{Y}_n;\mathcal{Y}_{0,n-1}\times\mathcal{X}_{0,n}):n\in\mathbb{N}\Big\}$, and the alternative directed information expression (III.16). These alternative formulations are described in the following remark.

Remark IV.2.

Part A. *For any arbitrary stochastic kernel $\bar{\nu}_{i|i-1}(dy_i;y^{i-1})\in\mathcal{Q}(\mathcal{Y}_i;\mathcal{Y}_{0,i-1})$, $i=0,1,\ldots,n$, then (IV.5) can be written as*

$$\begin{aligned} I(X^n\to Y^n) &\equiv \mathbb{I}_{X^n\to Y^n}(p_i(\cdot;\cdot,\cdot),q_i(\cdot;\cdot,\cdot):\ i=0,1,\ldots,n) \\ &= \inf_{\bar{\nu}_{i|i-1}(\cdot;\cdot)\in\mathcal{Q}(\mathcal{Y}_i;\mathcal{Y}_{0,i-1}):i=0,1,\ldots,n} \sum_{i=0}^n \int_{\mathcal{X}_{0,i}\times\mathcal{Y}_{0,i-1}} \mathbb{D}\big(q_i(\cdot;\cdot,\cdot)||\bar{\nu}_{i|i-1}(\cdot;\cdot)\big)(y^{i-1},x^i) \\ &\quad \times p_i(dx_i;x^{i-1},y^{i-1})\otimes P_{0,i-1}(dy^{i-1},dx^{i-1}) \end{aligned} \quad \text{(IV.17)}$$

where the infimum is achieved by $\bar{\nu}^*_{i|i-1}(dy_i;y^{i-1}) = \nu_{i|i-1}(dy_i;y^{i-1}),\ i=0,1,\ldots,n$. *The derivation is similar to the one of Theorem IV.1,* **Part A.**, *but it is done with respect to component* $\bar{\nu}_{i|i-1}$.

Part B. *For any arbitrary family of stochastic kernel* $\{r_i(\cdot;\cdot,\cdot)\in\mathcal{Q}(\mathcal{X}_i;\mathcal{X}_{0,i-1}\times\mathcal{Y}_{0,i-1}),\ i=0,1,\ldots,n\}$, *and* $\{s_i(\cdot;\cdot,\cdot)\in\mathcal{Q}(\mathcal{Y}_i;\mathcal{Y}_{0,i-1}\times\mathcal{X}_{0,i-1}),\ i=0,1,\ldots,n\}$, *define*

$$\mathbb{I}(p_i,q_i,s_i\otimes r_i:\ i=0,1,\ldots,n)\triangleq\sum_{i=0}^{n}\int_{\mathcal{X}_{0,i}\times\mathcal{Y}_{0,i}}\log\left(\frac{s_i(dy_i;y^{i-1},x^{i-1})\otimes r_i(dx_i;x^{i-1},y^i)}{p_i(dx_i;x^{i-1},y^{i-1})\otimes\nu_{i|i-1}(dy_i;y^{i-1})}\right)$$
$$\otimes_{k=0}^{i}p_k(dx_k;x^{k-1},y^{k-1})\otimes q_k(dy_k;y^{k-1},x^k).$$

Then (IV.7) can be written as

$$I(X^n\to Y^n)\equiv\mathbb{I}_{X^n\to Y^n}(p_i(\cdot;\cdot,\cdot),q_i(\cdot;\cdot,\cdot):\ i=0,1,\ldots,n)$$
$$=\sup_{\substack{s_i(\cdot;y^{i-1}x^{i-1})\otimes r_i(\cdot;x^{i-1}y^i):\ i=0,1,\ldots,n\\ s_i(\cdot;\cdot,\cdot)\in\mathcal{Q}(\mathcal{Y}_i;\mathcal{Y}_{0,i-1}\times\mathcal{X}_{0,i-1}),r_i(\cdot;\cdot,\cdot)\in\mathcal{Q}(\mathcal{X}_i;\mathcal{X}_{0,i-1}\times\mathcal{Y}_{0,i})}}\mathbb{I}(p_i,q_i,s_i\otimes r_i:\ i=0,1,\ldots,n) \tag{IV.18}$$

where the supremum is achieved by (IV.9) and (IV.11). The derivation is similar to the one of Theorem IV.1, **Part B.**, *but it is done with respect to component* $s_i\otimes r_i$, $i=0,1,\ldots,n$.

V. Conclusion

In this paper we derive functional and topological properties of directed information defined on abstract Polish spaces. We establish convexity of the sets of (nonanticipative) causally conditioned convolutional distributions, used to defined directed information and we show convexity and concavity of directed information with respect to these distributions. We provide a novel theorem on weak compactness of causally conditioned convolutional distributions, and weak convergence of joint distributions and marginal distributions associated with directed information. We use these results to show lower semicontinuity of directed information as a functional of two causally conditioned convolutional distributions, and under certain conditions continuity of directed information with respect to the causally conditioned convolutional input distributions. We also establish two variational equalities for directed information. These functional and topological properties may be viewed as generalizations of analogous functional and topological properties of mutual information. The results of this paper can be used to address extremum problems of directed information functional for point-to-point communications and for network communications.

VI. Appendix A

A. Backround material

In this section, we introduce the basic analytical concepts which are used throughout the paper.

Weak Convergence and Compactness.

The main notions discussed are weak convergence of probability measures, the relation to convergence with respect to Prohorov metric, tightness of a family of probability measures, relative compactness, weak compactness, and compactness.

Let $(\mathcal{X}, d)$ be a metric space, $\mathcal{B}(\mathcal{X})$ the $\sigma-$algebra of Borel subsets of $\mathcal{X}$, and $\mathcal{M}_1(\mathcal{X})$ is the family of probability measures on $\mathcal{X}$. Let $BC(\mathcal{X})$ denote the set of bounded, continuous real-valued function f on $(\mathcal{X}, d)$ endowed with the supremum norm $||f|| = \sup_{x\in\mathcal{X}} |f(x)|$. From [32] it is known that each $P \in \mathcal{M}_1(\mathcal{X})$ is uniquely determined by the integrals $\{\int_{\mathcal{X}} f(x)dP(x) : f \in BC(\mathcal{X})\}$. A sequence $\{P_n : n \geq 1\} \subset \mathcal{M}_1(\mathcal{X})$ is said to *converge weakly* to $P \in \mathcal{M}_1(\mathcal{X})$ if

$$\lim_{n\to\infty} \int_{\mathcal{X}} f(x)dP_n(x) = \int_{\mathcal{X}} f(x)dP(x), \ \forall f \in BC(\mathcal{X}).$$

Weak convergence of $\{P_n : n \geq 1\}$ to P is denoted by $P_n \xrightarrow{w} P$.

The space $\mathcal{M}_1(\mathcal{X})$ can be made into a Hausdorff topological space by taking basic neighbourhoods of $P \in \mathcal{M}_1(\mathcal{X})$ all sets the form

$$\left\{ Q \in \mathcal{M}_1(\mathcal{X}) : \Big| \int_{\mathcal{X}} f_i(x)dQ(x) - \int_{\mathcal{X}} f_i(x)dP(x) \Big| < \epsilon, i = 1, \ldots, k \right\}$$

where $\epsilon > 0$ and $f_1, f_2, \ldots, f_k \in BC(\mathcal{X})$. The resulting topology is called the *topology of weak convergence* or *weak topology*. Hence, a sequence $\{P_n : n \geq 1\}$ converges to P in this topology if and only if $P_n \xrightarrow{w} P$. The space $\mathcal{M}_1(\mathcal{X})$ is metrizable with respect to the Prohorov metric [40], denoted by $\mathcal{L}(\cdot, \cdot)$, and with respect to this metric $\mathcal{M}_1(\mathcal{X})$ is a Hausdorff topological space. The following theorem gives fundamental results regarding $(\mathcal{M}_1(\mathcal{X}), \mathcal{L})$.

Theorem VI.1. *[41, pp. 43–46]*

Let $(\mathcal{X}, d)$ be a metric space.

1) If $\mathcal{X}$ is separable, then $\mathcal{M}_1(\mathcal{X})$ is separable.

2) If $(\mathcal{X}, d)$ is complete and separable, then $(\mathcal{M}_1(\mathcal{X}), \mathcal{L})$ is a complete separable metric space (with respect to the Prohorov metric $\mathcal{L}$).

3) If $(\mathcal{X}, d)$ is compact metric space, then $(\mathcal{M}_1(\mathcal{X}), \mathcal{L})$ is a compact metric space.

Thus, separability is a topological property while completeness is a property of the metric. Statement Theorem VI.1, 3) (due to Prohorov) states that for $(\mathcal{X}, d)$ a metric space and $\mathcal{X}$ compact, then any sequence $\{P_n : n \geq 1\}$ of probability measures of $\mathcal{X}$ possess a convergent subsequence (with respect to the Prohorov metric).

A crucial result for the characterization of compact subsets of $\mathcal{M}_1(\mathcal{X})$ is the next theorem due to Prohorov, which relates compactness and tightness of a set of measures.

Definition VI.2. (*Tight Measures*) *[40, p. 308]*

A probability measure $P \in \mathcal{M}_1(\mathcal{X})$ is said to be tight if for every $\epsilon > 0$ there exists a compact set $K \subset \mathcal{X}$ such that $P(K) \geq 1 - \epsilon$ or $P(K^c) < \epsilon$. Moreover, a family of probability measures $M \subset \mathcal{M}_1(\mathcal{X})$ is said to be tight or uniformly tight if for every $\epsilon > 0$ there exists a compact set $K \subset \mathcal{X}$ such that $\inf_{P \in M} P(K) \geq 1 - \epsilon$.

Thus, $M \stackrel{\triangle}{=} \{P_n : n \geq 1\}$ is uniformly tight if for every $\epsilon > 0$ there exist a compact set $K \subset \mathcal{X}$ such that $P_n(K) \geq 1 - \epsilon, \ n = 1, 2, \ldots$. The definition of relative and weak compactness is defined below.

Definition VI.3. *(Relative and Weak Compactness)*

1) [32, Ch. 1, p. 57] A family of probability measures $M \subset \mathcal{M}_1(\mathcal{X})$ is said to be relatively compact (with respect to Prohorov metric) if each sequence $\{P_n : n \geq 1\}$ of elements in M contains some subsequence $\{P_{n_i} : i \in \{1, 2, \ldots\}\}$ converging (as $i \longrightarrow \infty$) to some probability measure $P \in \mathcal{M}_1(\mathcal{X})$ with respect to the Prohorov metric $\mathcal{L}(\cdot, \cdot)$. Here, the limit P is not required to belong to M, but all is required is to belong to $\mathcal{M}_1(\mathcal{X})$.

2) [32, Ch. 1, Corollary, p. 59] A family of probability measures $M \subset \mathcal{M}_1(\mathcal{X})$ is said to be weakly compact or relatively compact (with respect to weak convergence) if 1) holds with convergence tested with respect to weak convergence.

The following theorem due to Prohorov, relates weak convergence and tightness of a family of probability measures.

Theorem VI.4. (*Prohorov's Theorem*) *[40, Theorem A.3.15, p. 309]*
Let $(\mathcal{X}, d)$ be complete and separable metric space and let a family of probability measures M such that $M \subset \mathcal{M}_1(\mathcal{X})$. Then, the family of probability measures M is relatively compact with respect to weak convergence if and only if it is tight.

Note that if $(\mathcal{X}, d)$ is an arbitrary metric space and $M \subset \mathcal{M}_1(\mathcal{X})$ a family of probability measures, then tightness of M implies relative compactness of M, but the reverse is not valid. However, when $(\mathcal{X}, d)$ is complete separable metric space, then relative compactness and weak compactness are equivalent notions. Therefore, a family of probability measures $M \subset \mathcal{M}_1(\mathcal{X})$ on a complete separable metric space $(\mathcal{X}, d)$ is weakly compact or relatively compact with respect to weak convergence if and only if it is tight. Moreover, if $P_n \xrightarrow{w} P$, then the family $\{P_n : n \geq 1\}$ is tight.
Finally, we give another version due to Prohorov for a set of measures $M \subset \mathcal{M}_1(\mathcal{X})$ to be compact.

Theorem VI.5. *(Corollary of Prohorov's Theorem)*
Let $(\mathcal{X}, d)$ be a separable metric and $M \subset \mathcal{M}_1(\mathcal{X})$ a set of measures. The following hold.
(a) If M is closed and tight, then M is compact.
(b) Suppose $\mathcal{X}$ is complete. If M is compact then M is closed and tight.

That is, for $(\mathcal{X}, d)$ a separable metric space a sufficient condition for compactness of $M \subset \mathcal{M}_1(\mathcal{X})$ is that M is closed and tight. If $\mathcal{X}$ is also complete, then this condition is also necessary.

Lebesgue's Dominated Convergence Theorems (LDCT).
In this paper we often need to establish convergence of a sequence of integrals. Sufficient conditions are given by LDCT, which allow one to interchange the limit and the integral. We state these Lebesgue's theorems below.

Theorem VI.6. (*Lebesgue's Dominated Convergence Theorem (LDCT)*) *[31, Theorem 3, p. 187]*
Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $Y, X, X_1, X_2, \ldots$ be RV's such that $|X_n| \leq Y$, for all $n \geq 1$, $\mathbb{E}Y < \infty$ and $X_n \xrightarrow{a.s.} X$.
Then $\mathbb{E}|X| < \infty$,

a) $\lim_{n\to\infty} \mathbb{E}X_n = \mathbb{E}X$;

b) $\lim_{n\to\infty}\mathbb{E}|X_n - X| = 0$.

The previous theorem can be generalized as follows.

Corollary VI.7. *[31, Corollary, p. 188](LDCT in L^p-Spaces)*
Let $(\Omega,\mathcal{F},\mathbb{P})$ be a probability space and $Y, X, X_1, X_2, \ldots$ be RV's such that $|X_n| \leq Y$, $X_n \overset{a.s.}{\Longrightarrow} X$ and $\mathbb{E}Y^p < \infty$ for some $p > 0$. Then $\mathbb{E}|X|^p < \infty$ and $\lim_{n\to\infty}\mathbb{E}|X - X_n|^p = 0$.

Theorem VI.8. (*Generalized LDCT*) *[42, Exercise 20, p. 59]*
Let $\{f_n : n \geq 1\}$, $\{g_n : n \geq 1\}$, $f, g \in L^1$, $f_n \overset{a.s.}{\Longrightarrow} f$, $g_n \overset{a.s.}{\Longrightarrow} g$, $|f_n| \leq g_n$ for all $n \geq 1$, and $\lim_{n\to\infty}\int g_n d\mathbb{P} = \int g d\mathbb{P}$. Then $\lim_{n\to\infty}\int f_n d\mathbb{P} = \int f d\mathbb{P}$.

Uniform Integrability.
In this paper we shall also need stronger sufficient conditions to verify convergence of a sequence of integrals using the concept of uniform integrability. We state this next.

Definition VI.9. (*Uniform Integrability of RV's*) *[31, Definition 4, p. 188]*
Let $(\Omega,\mathcal{F},\mathbb{P})$ be a probability space. A sequence of RV's $\{X_n : n \geq 1\}$ is said to be uniformly $\mathbb{P}$-integrable if

$$\limsup_{c\to 0}\sup_{n\geq 1}\int_{\{\omega:|X_n(\omega)|\geq c\}}|X_n(\omega)|d\mathbb{P}(\omega) = 0.$$

Equivalently,

$$\sup_{n\geq 1}\mathbb{E}\Big\{|X_n|I_{\{|X_n)|>c\}}\Big\} \longrightarrow 0, \text{ as } c \to 0.$$

Note that if $\{X_n : n \geq 1\}$ satisfy $|X_n| \leq Y$ and $\mathbb{E}\{Y\} < \infty$, then the sequence $\{X_n : n \geq 1\}$ is uniformly integrable. Also, for the space $L^p(\Omega,\mathcal{F},\mathbb{P}), p \geq 1$, (the space of RV's with finite absolute $p-$th moments, $\mathbb{E}\big\{|X|^p\big\} < \infty$), if $\mathcal{L} \subset L^p(\Omega,\mathcal{F},\mathbb{P})$ is bounded for some $p > 1$ then $\mathcal{L}$ is uniformly integrable.
The following theorem gives some properties for a family of uniformly integrable RV's.

Theorem VI.10. (*Uniform Integrability of RV's*) *[31, Theorem 4, pp. 188-189]*
Let $(\Omega,\mathcal{F},\mathbb{P})$ be a probability space and $\{X_n : n \geq 1\}$ a uniformly $\mathbb{P}$-integrable family of RV's. Then

(a) $\mathbb{E}\liminf_n X_n \leq \liminf_n \mathbb{E}X_n \leq \limsup_n \mathbb{E}X_n \leq \mathbb{E}\limsup_n X_n$.

(b) *If in addition* $X_n \stackrel{a.s.}{\Longrightarrow} X$, *then* $\mathbb{E}|X| < \infty$, $\lim_{n\to\infty} \mathbb{E}|X_n| = \mathbb{E}|X|$ *and* $\lim_{n\to\infty} \mathbb{E}\Big\{|X_n - X|\Big\} = 0$.

Definition VI.9 describes uniform integrability when the integrand is a sequence and the probability measure is fixed. In the next definition of uniform integrability the integrand is fixed but the probability measure is a sequence.

Definition VI.11. (*Uniform Integrability for a family of probability measures*) *[36, Appendix, Definition A.2, p. 3084]*
Let $M \subset \mathcal{M}_1(\mathcal{X})$ *be a family of probability measures on* $\big(\mathcal{X}, \mathcal{B}(\mathcal{X})\big)$. *A measurable function* f *is said to be uniformly integrable over* M *if*

$$\lim_{c\to\infty} \sup_{P\in M} \int_{\{x\in\mathcal{X}:|f(x)|>c\}} |f(x)| dP(x) \longrightarrow 0.$$

A sufficient condition for the convergence of a sequence of integrals of a function with respect to a weakly convergent sequence of measures is the following.

Theorem VI.12. *[36, Appendix, Theorem A.2, p. 3084]*
Let $M \subset \mathcal{M}_1(\mathcal{X})$ *be a closed family of probability measures on* $\big(\mathcal{X}, \mathcal{B}(\mathcal{X})\big)$, *and let* $\{P_n : n \geq 1\} \subset M$ *be a weakly convergent sequence in* M. *If* f *is a continuous function and uniformly integrable over* $\{P_n : n \geq 1\}$ *then* $\lim_{n\to\infty} \int f dP_n = \int f dP$.

Regular Conditional Probability Measures and their Properties.
In this paper we extensively utilized conditional distribution and expectation. Hence we shall need their precise definitions, given in terms of the Radon-Nikodym Derivative.
For any two measurable spaces $(E, \mathcal{E})$ and $(F, \mathcal{F})$, denote by $\mathcal{E} \odot \mathcal{F}$, the σ-algebra on $E \times F$ generated by the collection of all measurable rectangles $A \times B \stackrel{\triangle}{=} \{(x, y) : x \in A, y \in B\}$, $A \subset \mathcal{E}$, $B \subset \mathcal{F}$, called the product σ-algebra; the measurable space $(E \times F, \mathcal{E} \odot \mathcal{F})$ is called the product space of $(E, \mathcal{E})$ and $(F, \mathcal{F})$.
Let $(\Omega, \mathcal{F})$ be a measurable space. Given two probability measures P, Q on $(\Omega, \mathcal{F})$, Q is said to be *absolutely continuous* with respect to P (denoted $P \ll Q$) if for every $A \in \mathcal{F}$ such that $P(A) = 0$ then $Q(A) = 0$. If $Q \ll P$, by Radon-Nikodym Derivative theorem, there exists a $P-$integrable and $\mathcal{F}-$measurable function f such that for every $A \in \mathcal{F}$, $Q(A) = \int_A f(\omega) dP(\omega)$. The function f is unique $P - a.e.$ (almost everywhere) and is called the *Radon-*

Nikodym Derivative (RND) density of Q with respect to P, denoted by $f(\omega) = \frac{dQ}{dP}|_{\mathcal{F}(\omega)}$. That is, if $\bar{f}$ is another function satisfying these properties, there exists a $P-$null set N such that $f(\omega) = \bar{f}(\omega),\ \forall \omega \in N^c \stackrel{\triangle}{=} (\Omega \setminus N)$.This relation is denoted by $f(\omega) = \bar{f}(\omega),\ P - a.s.$, and we say that the RND f is unique $P - a.s.$ If in addition to $Q \ll P$, also $P \ll Q$ holds, then P and Q are called *equivalent* denoted by $Q \sim P$.

Next, we state the definition of a regular conditional probability measure, from which the regular conditional distribution of one RV given another RV can be defined.

Definition VI.13. (*Regular Conditional Probability Measure*) *[40, A.4, pp. 312-313]*

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\mathcal{G}$ be a sub-σ-algebra of $\mathcal{F}$. A regular conditional probability measure $P(\cdot|\mathcal{G})(\cdot)$ on $(\Omega, \mathcal{F})$ is a function $P(A|\mathcal{G})(\omega),\ A \in \mathcal{F},\ \omega \in \Omega$ having the following properties.

(a) For each $A \in \mathcal{F}$, the function mapping $\omega \in \Omega \longmapsto P(A|\mathcal{G})(\omega)$ is measurable with respect to $\mathcal{G}$.

(b) For each $\omega \in \Omega$, $P(\cdot|\mathcal{G})(\omega)$ is a probability measure on $\mathcal{F}$.

(c) For each $A \in \mathcal{F}$, $P(A|\mathcal{G})(\omega)$ is a version of the conditional probability of A given $\mathcal{G}$. Moreover,

$$P(A \cap B) = \int_B P(A|\mathcal{G})(\omega) P_{\mathcal{G}}(d\omega),\ \forall A \in \mathcal{G}$$

where $P_{\mathcal{G}}$ is the restriction of P to $\mathcal{G}$.

Statements (a) and (c) state that $P(A|\mathcal{G})(\omega)$ is a version of the conditional probability of A given $\mathcal{G}$ (and it is a function of ω). If such a version $P(\cdot|\mathcal{G})(\cdot)$ exists then it is unique in the sense that, if $\bar{P}(\cdot|\mathcal{G})(\cdot)$ is another function with these properties, then there exists a $P_{\mathcal{G}}-$null set N such that $P(A|\mathcal{G})(\omega) = \bar{P}(A|\mathcal{G})(\omega),\ \forall A \in \mathcal{F}$ and $\omega \in N^c$ (e.g., $P(\cdot|\mathcal{G})(\omega) = \bar{P}(\cdot|\mathcal{G})(\omega),\ P_{\mathcal{G}} - a.s.$). Thus, a regular conditional probability measure exists if it can be shown that a version of the conditional probability measure can be chosen to be a probability measure on $\mathcal{F}$ for each $\omega \in \Omega$. Although in general, a regular conditional probability measure may not exist, for the case when $\mathcal{G}$ is generated by a countable partition of Ω, a regular conditional probability measure given $\mathcal{G}$ always exists. Moreover, if (Ω, d) is a metric space which is complete and separable (Polish space), and $\mathcal{F}$ is a Borel $\sigma-$algebra, then for any probability measure P on $(\Omega, \mathcal{F})$ and any sub-σ-algebra $\mathcal{G} \subseteq \mathcal{F}$, a regular conditional probability measure of P given $\mathcal{G}$ always exists.

The next lemma summarizes certain relationships between the absolute continuity of probability measures.

Lemma VI.14. *(Absolute Continuity of Probability Measures) [43, Lemma 4.4.7, pp. 149-150]*
a) Suppose $Q_{\mathcal{G}} \ll P_{\mathcal{G}}$. *If* $Q(\cdot|\mathcal{G})(\omega) \ll P(\cdot|\mathcal{G})(\omega),\ Q_{\mathcal{G}} - a.s.$, *then* $Q \ll P$.
b) Conversely, if $Q \ll P$, *then* $Q(\cdot|\mathcal{G})(\omega) \ll P(\cdot|\mathcal{G})(\omega),\ P(\cdot|\mathcal{G})(\omega) - a.s.$

Note that if $Y : (\Omega, \mathcal{F}) \longmapsto (\mathcal{Y}, \mathcal{A})$ is a RV on $(\Omega, \mathcal{F})$ into a measurable space $(\mathcal{Y}, \mathcal{A})$ and $\mathcal{Y}$ is a Polish space, then a regular conditional distribution for Y given the sub-σ-algebra $\mathcal{G}$ of $\mathcal{F}$ denoted by $P(dy|\mathcal{G})(\omega)$ is defined according to Definition VI.13, and this always exists.
Additionally, if $X : (\Omega, \mathcal{F}) \longmapsto (\mathcal{X}, \mathcal{B})$ is a RV on $(\Omega, \mathcal{F})$ into a measurable space $(\mathcal{X}, \mathcal{B})$, and $\mathcal{G}$ is the sub-σ-algebra of $\mathcal{F}$ generated by X, then $P(dy|X)(\omega)$ is called the *regular conditional distribution* of Y given X. One can go one step further to define a regular conditional distribution for Y given $X = x$ as a quantity $P(dy|X = x)$, and introduce an equivalent definition called stochastic kernel.

Definition VI.15. (*Stochastic Kernel*) *[40, p. 28 and Theorem A.5.2, pp. 316-317]*
Consider the measurable spaces $(\mathcal{X}, \mathcal{B}), (\mathcal{Y}, \mathcal{A})$.
A stochastic Kernels is a mapping $q : \mathcal{A} \times \mathcal{X} \to [0, 1]$ *satisfying the following two properties:*
1) For every $x \in \mathcal{X}$, *the set function* $q(\cdot; x)$ *is a probability measure on* $\mathcal{A}$;
2) for every $F \in \mathcal{A}$, *the function* $q(F; \cdot)$ *is* $\mathcal{B}$*-measurable.*
The set of stochastic kernels on $\mathcal{Y}$ *given* $\mathcal{X}$ *is denoted by* $\mathcal{Q}(\mathcal{Y}; \mathcal{X})$.

The next important result relates a certain measurability called consistency of conditional distributions to the existence of a sequence of stochastic kernels.

Theorem VI.16. *(Equivalence of Consistent Family of Measures and a Sequence of Conditional Distributions) [21, Chapter 1, Theorem 1.1, pp. 4-5]*
Let $\{(\mathcal{X}_n, \mathcal{B}(\mathcal{X}_n)) : n \in \mathbb{N}\}$ *be complete separable metric spaces (Polish spaces) with* $\mathcal{B}(\mathcal{X}_n)$ *a* $\sigma-$*algebra of Borel sets. For any family of measures* $\mathbf{P}(\cdot|\mathbf{y})$ *on* $(\mathcal{X}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}))$ *satisfying consistency condition*
C1*: If* $E \in \mathcal{B}(\mathcal{X}_{0,n})$ *then* $\mathbf{P}(E_{0,n}|\mathbf{y})$ *is* $\mathcal{B}(\mathcal{Y}_{0,n-1})-$*measurable function of* $\mathbf{y} \in \mathcal{Y}^{\mathbb{N}}$
there exists a family of conditional distributions $\{p_n(dx_n; x^{n-1}, y^{n-1}) : n \in \mathbb{N}\}$ *satisfying*

conditions

i) *For every* $n \in \mathbb{N}$, $p_n(\cdot; x^{n-1}, y^{n-1})$ *is a probability measure on* $\mathcal{B}(\mathcal{X}_n)$*;*

ii) *For every* $n \in \mathbb{N}$, $p_n(A_n; x^{n-1}, y^{n-1})$ *is* $\odot_{i=0}^{n-1}\big(\mathcal{B}(\mathcal{X}_i) \odot \mathcal{B}(\mathcal{Y}_i)\big)$*-measurable function of* $x^{n-1} \in \mathcal{X}_{0,n-1}$, $y^{n-1} \in \mathcal{Y}_{0,n-1}$*;*

such that

$$\mathbf{P}(C|\mathbf{y}) \stackrel{\triangle}{=} \int_{C_0} p_0(dx_0) \int_{C_1} p_1(dx_1; x^0, y^0) \ldots \int_{C_n} p_n(dx_n; x^{n-1}, y^{n-1}),\ C = \times_{i=0}^{n} C_i.$$

Convexity of Regular Conditional Distributions.

Next, we give the definition of the well-known convexity properties of the family of conditional distributions.

Let $\mathcal{M}_1(\Omega, \mathcal{F}, \mathcal{G}, P),\ \mathcal{G} \subseteq \mathcal{F}$ denote the set of all probability measures Q on $(\Omega, \mathcal{F})$ such that $Q \ll P$ and $Q_{\mathcal{G}} \sim P_{\mathcal{G}}$. Here P is a fixed probability measure. Let $\mathcal{M}_1(\Omega, \mathcal{F}, P|\mathcal{G})(\omega)$ denote the set of all regular conditional probability measures on $(\Omega, \mathcal{F})$ conditional on $\mathcal{G}$ defined by $\mathcal{M}_1(\Omega, \mathcal{F}, P|\mathcal{G})(\omega) \stackrel{\triangle}{=} \{Q(\cdot|\mathcal{G}) : Q \in \mathcal{M}_1(\Omega, \mathcal{F}, \mathcal{G}, P)\}$. Next, we define the almost sure-convexity of the set of regular conditional distributions.

Definition VI.17. *(Almost Sure Convexity of a Set of Regular Conditional Probability Measures) We say that the set* $\mathcal{M}_1(\Omega, \mathcal{F}, P|\mathcal{G})(\omega)$ *is convex if the following property holds.*

Given $Q_1(\cdot|\mathcal{G})(\omega)$, $Q_2(\cdot|\mathcal{G})(\omega)$ *in* $\mathcal{M}_1(\Omega, \mathcal{F}, P|\mathcal{G})(\omega)$ *and* $\lambda \in (0, 1)$*, there exists a probability measure* $\hat{Q}$ *on* $(\Omega, \mathcal{F})$ *whose regular conditional probability measure* $\hat{Q}(\cdot|\mathcal{G}(\omega) \in \mathcal{M}_1(\Omega, \mathcal{F}, P|\mathcal{G})(\omega)$ *satisfies* $\hat{Q}(\cdot|\mathcal{G})(\omega) = Q_1(\cdot|\mathcal{G})(\omega) + (1-\lambda)Q_2(\cdot|\mathcal{G})(\omega),\ P_{\mathcal{G}}$*-a.s. That is, there exists a* $P_{\mathcal{G}}$*-null set* N *such that for such* $D \in \mathcal{F}$ *and for all* $\omega \in N^c$,

$$\hat{Q}(D|\mathcal{G})(\omega) = Q_1(D|\mathcal{G})(\omega) + (1-\lambda)Q_2(D|\mathcal{G})(\omega).$$

This property is denoted by $\lambda Q_1(\cdot|\mathcal{G})(\omega) + (1-\lambda)Q_2(\cdot|\mathcal{G})(\omega) \in \mathcal{M}_1(\Omega, \mathcal{F}, P|\mathcal{G})(\omega)$.

Relative Entropy.

Mutual information is often defined via the relative entropy or Kullback-Leibler distance between two distributions. Next, we give the definition and the chain rule of relative entropy.

Definition VI.18. (*Relative Entropy*) *[30, Definition 1.4.1, p. 21]*

Given a measurable space $(\mathcal{Z}, \mathcal{B}(\mathcal{Z}))$*, the relative entropy between two probability measures*

$P \in \mathcal{M}_1(\mathcal{Z})$ and $Q \in \mathcal{M}_1(\mathcal{Z})$ is defined by

$$\mathbb{D}(P||Q) \stackrel{\triangle}{=} \begin{cases} \int_{\mathcal{Z}} \log\left(\frac{dP}{dQ}\right) dP = \int_{\mathcal{Z}} \log\left(\frac{dP}{dQ}\right) \frac{dP}{dQ} dQ & \text{if} \quad P << Q \\ +\infty & \text{otherwise} \end{cases}$$

where $\frac{dP}{dQ}$ denotes the RND (density) of P with respect to Q, and $P << Q$ denotes absolute continuity of Q with respect to P (e.g., $Q(A) = 0$ for some $A \in \mathcal{B}(\mathcal{Z})$ then $P(A) = 0$).

The previous definition is often used to define mutual information between RV's X and Y via $I(X;Y) \stackrel{\triangle}{=} \mathbb{D}(P_{X,Y}||P_X \times P_Y)$. However, it is often desirable to express $I(X;Y)$ as a functional of distribution P_X and conditional distribution $P_{Y|X}$. This is established via the chain rule of relative entropy given below.

Theorem VI.19. (*Chain Rule of Relative Entropy*) *[40, Theorem B.2.1, p. 326]*
Let $(\mathcal{X}, \mathcal{B})$ and $(\mathcal{Y}, \mathcal{A})$ be polish (complete separable metric spaces) spaces and P and Q be probability measures on $\mathcal{X} \times \mathcal{Y}$. Let P_1 and Q_1 the first marginals of P and Q $(e.g., P_1(A) = P(A \times \mathcal{Y})$ and $Q_1(A) = Q(A \times \mathcal{Y}), \forall A \in \mathcal{B})$ respectively, and by $\alpha(dy;x)$ and $\beta(dy;x)$ the stochastic kernels on $\mathcal{Y}$ given $\mathcal{X}$ for which we have the decomposition, $P(dx,dy) = P_1(dx) \otimes \alpha(dy;x)$ and $Q(dx,dy) = Q_1(dx) \otimes \beta(dy;x)$. Then the function mapping $x \mapsto H(\alpha(\cdot;x)||\beta(\cdot;x))$ is measurable and

$$H(P||Q) = H(P_1||Q_1) + \int_{\mathcal{X}} H(\alpha(\cdot;x)||\beta(\cdot;x)) P_1(dx).$$

By invoking the chain rule of relative entropy with an appropriate choice of P and Q and Lemma VI.14, it can be shown that $I(X;Y) = \int_{\mathcal{X}} \mathbb{D}\big(P_{Y|X}(\cdot|x)||P_Y(\cdot)\big) \times P_X(dx)$.

VII. Appendix B

A. Additional Discussion

1) Directed Information as a Functional of Sequence of Conditional Distributions: The equivalent representations (III.10)-(III.12) are obtained from the definition of conditional mutual information $I(X^i; Y_i|Y^{i-1})$, by invoking the chain rule of relative entropy and the relation between absolute continuity of measures (Appendix VI-A, Theorem VI.19 and Lemma VI.14). These are also discussed formally in [40]. Indeed, if $P_{0,i}(\cdot,\cdot;y^{i-1}) \ll P_{0,i}(\cdot;y^{i-1}) \times \nu_{i|i-1}(\cdot;y^{i-1})$, $\nu_{i-1}-$almost all $y^{i-1} \in \mathcal{Y}_{0,i-1}$ then by the Radon-Nikodym theorem there exists a version of

the RND $\xi_i(x^i,y^i) \stackrel{\triangle}{=} \frac{P_{0,i}(\cdot,\cdot;y^{i-1})}{P_{0,i}(\cdot;y^{i-1})\times\nu_{i|i-1}(\cdot;y^{i-1})}(x^i,y_i)$ which is non-negative for all $i=0,1,\ldots,n$. Moreover, by Lemma VI.14, $P_{0,i}(\cdot,\cdot;y^{i-1}) \ll P_{0,i}(\cdot;y^{i-1})\times\nu_{i|i-1}(\cdot;y^{i-1})$, $\nu_{0,i-1}-$almost all $y^{i-1}\in\mathcal{Y}_{0,i-1}$ if and only if $q_i(\cdot;y^{i-1},x^i)\ll\nu_{i|i-1}(\cdot;y^{i-1})$, $P_{0,i}-$almost all $(x^i,y^{i-1})\in\mathcal{X}_{0,i}\times\mathcal{Y}_{0,i-1},\ i=0,1,\ldots,n$, and in this case the RND $\xi_i(x^i,y^i)$ represents a version of $\frac{q_i(\cdot;y^{i-1},x^i)}{\nu_{i|i-1}(\cdot;y^{i-1})}(y_i)$, $P_{0,i}-$almost all $(x^i,y^{i-1})\in\mathcal{X}_{0,i}\times\mathcal{Y}_{0,i-1},\ i=0,1,\ldots,n$ (formally obtained via the decomposition $P_{0,i}(dx^i,dy_i;y^{i-1})=P_{0,i}(dx^i;y^{i-1})\otimes q_i(dy_i;y^{i-1},x^i)$).

Hence, when the RND $\xi_i(x^i,y^i)$ exists, repeated application of chain rule, Theorem VI.19 or following the derivation of Theorem in [40, Theorem B.2.1] (this is a very lengthy procedure), yields that (III.12) is obtained from (III.10). Similarly, starting with (III.12) one also obtains (III.10). Note that (III.10) and (III.12) are valid even when the RND do not exist in which case (III.10) and (III.12) take the value $+\infty$.

2) Directed Information as the supremum over finite partitions: In this section we illustrate how directed information can be equivalently defined as a supremum over appropriate partition of measurable spaces with respect to $\mathbf{P}(\cdot|\cdot)\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}};\mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\cdot)\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}};\mathcal{X}^{\mathbb{N}})$. Recall that relative entropy or information divergence of two probability measures $\mathbb{P}$ and $\mathbb{Q}$ (on a σ-algebra) of subsets of a Polish space is also defined [40] as

$$\mathbb{D}(\mathbb{P}||\mathbb{Q}) \stackrel{\triangle}{=} \sup_{\pi\in\Pi(\mathcal{Z})}\sum_{A\in\pi}\mathbb{P}(A)\log\frac{\mathbb{P}(A)}{\mathbb{Q}(A)} \tag{VII.1}$$

where $\Pi(\mathcal{Z})$ denotes the class of all finite measurable partitions of $\mathcal{Z}$. In (VII.1), the summation takes the value 0 if $\mathbb{P}(A)=0$ and the value $+\infty$ if $\mathbb{P}(A)>0$ and $\mathbb{Q}(A)=0$. Thus, $\pi\stackrel{\triangle}{=}\{A_1,\ldots,A_{m(\Pi)}\}\in\Pi(\mathcal{Z})$ is a partition of $\mathcal{Z}$ if A_j, $j=1,\ldots,m(\Pi)$, are measurable and $A_i\cap A_j=\{\emptyset\}$, $\forall i\neq j$, $\bigcup_{i=1}^{m(\Pi)}=\mathcal{Z}$. According to Dobrushin's theorem [29], under certain conditions mutual information $I(X;Y)=\mathbb{D}(P_{X,Y}||P_X\times P_Y)$ of two arbitrary RV's X and Y can be represented as a supremum of mutual information of discrete RV's corresponding to quantization of Y and X. Next, we utilize Dobrushin's theorem to represent directed information. Recall the following condition from Pinsker [29]. Given any set $\mathcal{Z}$ and a σ-algebra of its subsets, let $\Delta(\mathcal{Z})$ be a class of partitions of $\pi\stackrel{\triangle}{=}\{A_1,\ldots,A_{m(\Pi)}\}$ of $\mathcal{Z}$ such that 1) any $\pi_1\in\Delta(\mathcal{Z})$ and $\pi_2\in\Delta(\mathcal{Z})$ have a common refinement $\pi\in\Delta(\mathcal{Z})$, and 2) the algebra of all finite unions of atoms of partitions $\Delta(\mathcal{Z})$ generates the given σ-algebra. Then (VII.1) is valid with the supremum taken over $\pi\in\Delta(\mathcal{Z})\subseteq\Pi(\mathcal{Z})$.

For product spaces $\mathcal{Z}\stackrel{\triangle}{=}\mathcal{X}\times\mathcal{Y}$, 1) and 2) hold for the class of product partitions $\pi\stackrel{\triangle}{=}A\times B$ with

atoms $A_i \times B_j$, $i = 1, \ldots, m(A)$, $j = 1, \ldots, m(B)$, if A and B range over classes of partitions $\mathcal{X}$ and $\mathcal{Y}$, respectively, having properties 1) and 2).

Let $\mathcal{Z}_{0,n} \stackrel{\triangle}{=} \mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}$ and consider the class of product partitions $\pi_{0,n} \stackrel{\triangle}{=} \times_{k=0}^{n}(A_k \times B_k)$ with $A_k \times B_k$ atoms $A_{k,i} \times B_{k,j}$, $i = 1, \ldots, m(A_k)$, $j = 1, \ldots, m(B_k)$, where A_k and B_k range over the classes of partitions of $\mathcal{X}_k$ and $\mathcal{Y}_k$, respectively, having properties 1) and 2), $k = 0, 1, \ldots, n$, denoted by $\Delta(\mathcal{Z}_{0,n})$.

For any $\mathbf{P}(\cdot|\mathbf{y}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}^{\mathbb{N}}; \mathcal{Y}^{\mathbb{N}})$ and $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}^{\mathbb{N}}; \mathcal{X}^{\mathbb{N}})$, directed information is defined by

$$\mathbb{D}(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n} || \overrightarrow{\Pi}_{0,n}) \stackrel{\triangle}{=} \sup_{\pi_{0,n} \in \Delta(\mathcal{Z}_{0,n})} \sum_{A_{0,n} \in \pi_{0,n}} \log \frac{(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(A_{0,n})}{\overrightarrow{\Pi}_{0,n}(A_{0,n})} (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(A_{0,n}). \tag{VII.2}$$

Similarly, an equivalent definition of directed information as a supremum over partitions is obtained via (III.18).

B. Proof of Theorem III.4

1) Fix $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and let $\overrightarrow{Q}^1_{0,n}(\cdot|x^n)$, $\overrightarrow{Q}^2_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$. Then, the joint distributions corresponding to $\overrightarrow{Q}^1_{0,n}(\cdot|x^n)$, $\overrightarrow{Q}^2_{0,n}(\cdot|x^n)$ are

$$(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^1_{0,n})(dx^n, dy^n) \text{ and } (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^2_{0,n})(dx^n, dy^n),$$

and the marginals are

$$\nu^1_{0,n}(dy^n) = (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^1_{0,n})(\mathcal{X}_{0,n}, dy^n), \quad \nu^2_{0,n}(dy^n) = (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^2_{0,n})(\mathcal{X}_{0,n}, dy^n).$$

Since the set $\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$ is convex, given $\lambda \in (0, 1)$ there exists a probability measure $\tilde{\mathbf{P}}$ on $(\mathcal{X}^{\mathbb{N}} \times \mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}) \odot \mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ whose regular conditional measure $\mathbf{Q}(\cdot|\mathbf{x}) \in \mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}} \times \mathcal{Y}^{\mathbb{N}}, \mathcal{B}(\mathcal{X}^{\mathbb{N}}) \odot \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \mathcal{B}(\mathcal{Y}^{\mathbb{N}}), \bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{X}_{0,n})\big)(x^n)$ satisfies

$$\overrightarrow{Q}_{0,n}(\cdot|x^n) = \lambda \overrightarrow{Q}^1_{0,n}(\cdot|x^n) + (1-\lambda)\overrightarrow{Q}^2_{0,n}(\cdot|x^n), \ \bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{X}_{0,n})} - a.s.$$

and $\mathbf{C1}$ holds. Define

$$\nu_{0,n}(dy^n) = \lambda \nu^1_{0,n}(dy^n) + (1-\lambda)\nu^2_{0,n}(dy^n).$$

Introduce the RNDs $\Lambda^i_{0,n}(x^n,y^n)=\frac{\overrightarrow{Q}^i_{0,n}(dy^n|x^n)}{\nu^i_{0,n}(dy^n)}$, $\Psi^i_{0,n}(x^n,y^n)=\frac{\overrightarrow{Q}^i_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}$, $K^i_{0,n}(y^n)=\frac{\nu^i_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}$ and $\Lambda_{0,n}(x^n,y^n)=\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}$, $i=1,2$. Then,

$$\begin{aligned}\lambda\Psi^1_{0,n}(x^n,y^n)+(1-\lambda)\Psi^2_{0,n}(x^n,y^n)&=\lambda\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}+(1-\lambda)\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\\&=\frac{\lambda\overrightarrow{Q}^1_{0,n}(dy^n|x^n)+(1-\lambda)\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\lambda\nu^1_{0,n}(dy^n)+(1-\lambda)\nu^2_{0,n}(dy^n)}\\&=\Lambda_{0,n}(x^n,y^n)\end{aligned}$$

and

$$\begin{aligned}\lambda K^1_{0,n}(y^n)+(1-\lambda)K^2_{0,n}(y^n)&=\lambda\frac{\nu^1_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}+(1-\lambda)\frac{\nu^2_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}\\&=\frac{\lambda\nu^1_{0,n}(dy^n)+(1-\lambda)\nu^2_{0,n}(dy^n)}{\lambda\nu^1_{0,n}(dy^n)+(1-\lambda)\nu^2_{0,n}(dy^n)}=1.\end{aligned}$$

Applying the log-sum formula [44, Theorem 2.7.1, p. 31] yields

$$\begin{aligned}&\lambda\Psi^1_{0,n}(x^n,y^n)\log\Lambda^1_{0,n}(x^n,y^n)+(1-\lambda)\Psi^2_{0,n}(x^n,y^n)\log\Lambda^2_{0,n}(x^n,y^n)\\&=\lambda\Psi^1_{0,n}(x^n,y^n)\log\left(\frac{\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}}{\frac{\nu^1_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}}\right)+(1-\lambda)\Psi^2_{0,n}(x^n,y^n)\log\left(\frac{\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}}{\frac{\nu^2_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}}\right)\\&=\lambda\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\log\left(\frac{\lambda\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}}{\lambda\frac{\nu^1_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}}\right)+(1-\lambda)\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\log\left(\frac{(1-\lambda)\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}}{(1-\lambda)\frac{\nu^2_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}}\right)\\&\geq\left(\lambda\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}+(1-\lambda)\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\right)\log\left(\frac{\lambda\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}+(1-\lambda)\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}}{\lambda\frac{\nu^1_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}+(1-\lambda)\frac{\nu^2_{0,n}(dy^n)}{\nu_{0,n}(dy^n)}}\right)\\&=\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\log\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}.\end{aligned}$$

Integrating the above with respect to $\nu_{0,n}(dy^n)\otimes\overleftarrow{P}_{0,n}(dx^n|y^{n-1})$ yields:

$$\begin{aligned}&\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\big(\nu_{0,n}(dy^n)\otimes\overleftarrow{P}_{0,n}(dx^n|y^{n-1})\big)\\&=\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\end{aligned}$$

$$\leq \lambda \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu^1_{0,n}(dy^n)}\Big)(\overrightarrow{Q}^1_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)$$
$$+(1-\lambda)\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu^2_{0,n}(dy^n)}\Big)(\overrightarrow{Q}^2_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n).$$

Hence,

$$\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\lambda\overrightarrow{Q}^1_{0,n}+(1-\lambda)\overrightarrow{Q}^2_{0,n}\big)\leq\lambda\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\overrightarrow{Q}^1_{0,n}\big)+(1-\lambda)\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\overrightarrow{Q}^2_{0,n}\big).$$

This completes the derivation of 1).

2) Fix $\overrightarrow{Q}_{0,n}(\cdot|x^n)\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ and let $\overleftarrow{P}^1_{0,n}(\cdot|y^{n-1})$, $\overleftarrow{P}^2_{0,n}(\cdot|y^{n-1})\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. Then, the joint distributions corresponding to $\overleftarrow{P}^1_{0,n}(\cdot|y^{n-1})$, $\overleftarrow{P}^2_{0,n}(\cdot|y^{n-1})$ are

$$(\overleftarrow{P}^1_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n)\text{ and }(\overleftarrow{P}^2_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n).$$

The marginals corresponding to $\overleftarrow{P}^1_{0,n}(\cdot|y^{n-1})$, $\overleftarrow{P}^2_{0,n}(\cdot|y^{n-1})$ are

$$\nu^1_{0,n}(dy^n)=(\overleftarrow{P}^1_{0,n}\otimes\overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n},dy^n),\ \nu^2_{0,n}(dy^n)=(\overleftarrow{P}^2_{0,n}\otimes\overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n},dy^n).$$

Since the set $\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$ is convex, given $\lambda\in(0,1)$ there exists a probability measure $\tilde{\mathbf{P}}$ on $(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}},\mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}))$ whose regular conditional measure $\mathbf{P}(\cdot|\mathbf{y})\in\mathcal{M}_1\big(\mathcal{X}^{\mathbb{N}}\times\mathcal{Y}^{\mathbb{N}},\mathcal{B}(\mathcal{X}^{\mathbb{N}})\odot\mathcal{B}(\mathcal{Y}^{\mathbb{N}}),\mathcal{B}(\mathcal{X}^{\mathbb{N}}),\bar{\mathbf{P}}\big|\mathcal{B}(\mathcal{Y}_{0,n-1})\big)(y^{n-1})$ satisfies

$$\overleftarrow{P}_{0,n}(\cdot|y^{n-1})=\lambda\overleftarrow{P}^1_{0,n}(\cdot|y^{n-1})+(1-\lambda)\overleftarrow{P}^2_{0,n}(\cdot|y^{n-1}),\ \bar{\mathbf{P}}\big|_{\mathcal{B}(\mathcal{Y}_{0,n-1})}-a.s.$$

and $\mathbf{C2}$ holds. Then, corresponding to $\overleftarrow{P}_{0,n}(dx^n|y^{n-1})$ and $\overrightarrow{Q}_{0,n}(dy^n|x^n)$ we have

$$\begin{aligned}\nu_{0,n}(dy^n)&=\int_{\mathcal{X}_{0,n}}\big(\lambda\overleftarrow{P}^1_{0,n}(dx^n|y^{n-1})+(1-\lambda)\overleftarrow{P}^1_{0,n}(dx^n|y^{n-1})\big)\otimes\overrightarrow{Q}_{0,n}(dy^n|x^n)\\&=\lambda(\overleftarrow{P}^1_{0,n}\otimes\overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n},dy^n)+(1-\lambda)(\overleftarrow{P}^2_{0,n}\otimes\overrightarrow{Q}_{0,n})(\mathcal{X}_{0,n},dy^n)\\&=\lambda\nu^1_{0,n}(dy^n)+(1-\lambda)\nu^2_{0,n}(dy^n).\end{aligned}$$

Pick any measure $U_{0,n}(dy^n)\in\mathcal{M}_1(\mathcal{Y}_{0,n})$ with $\mathbb{D}(\nu_{0,n}||U_{0,n})<\infty$, e.g., such that $\nu_{0,n}(\cdot)\ll U_{0,n}(\cdot)$. Since $\overrightarrow{Q}(\cdot|x^n)\ll\nu_{0,n}(\cdot)$, for almost all $x^n\in\mathcal{X}_{0,n}$, and $\nu_{0,n}(\cdot)\ll U_{0,n}(\cdot)$, then $\overrightarrow{Q}_{0,n}(\cdot|x^n)\ll U_{0,n}(\cdot)$,

for almost all $x^n \in \mathcal{X}_{0,n}$. Consider

$$\begin{aligned}
\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}\big) &= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)\times U_{0,n}(dy^n)}{\nu_{0,n}(dy^n)\times U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&\quad - \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\nu_{0,n}(dy^n)}{U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&\quad - \int_{\mathcal{Y}_{0,n}} \log\Big(\frac{\nu_{0,n}(dy^n)}{U_{0,n}(dy^n)}\Big)\Big(\int_{\mathcal{X}_{0,n}}(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\Big)\\
&= \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&\quad - \int_{\mathcal{Y}_{0,n}} \log\Big(\frac{\nu_{0,n}(dy^n)}{U_{0,n}(dy^n)}\Big)\nu_{0,n}(dy^n).
\end{aligned}$$

Hence,

$$\begin{aligned}
&\mathbb{I}_{X^n\to Y^n}\big(\lambda\overleftarrow{P}^1_{0,n}+(1-\lambda)\overleftarrow{P}^2_{0,n}, \overrightarrow{Q}_{0,n}\big) = \int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}\Big)\\
&\times \overrightarrow{Q}_{0,n}(dy^n|x^n)\otimes\big(\lambda\overleftarrow{P}^1_{0,n}(dx^n|y^{n-1})+(1-\lambda)\overleftarrow{P}^2_{0,n}(dx^n|y^{n-1})\big) - \int_{\mathcal{Y}_{0,n}} \log\left(\frac{\nu_{0,n}(dy^n)}{U_{0,n}(dy^n)}\right)\nu_{0,n}(dy^n).
\end{aligned}$$

Moreover, relative entropy is convex in both arguments $\big($e.g., $\mathbb{D}(\cdot||U_{0,n})$ is convex for fixed $U_{0,n}\big)$, hence

$$\begin{aligned}
\mathbb{I}_{X^n\to Y^n}\big(\lambda\overleftarrow{P}^1_{0,n}+(1-\lambda)\overleftarrow{P}^2_{0,n}, \overrightarrow{Q}_{0,n}\big) &\geq \lambda\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}^1_{0,n})(dx^n,dy^n)\\
&\quad - \lambda\int_{\mathcal{Y}_{0,n}} \log\Big(\frac{\nu^1_{0,n}(dy^n)}{U_{0,n}(dy^n)}\Big)\nu^1_{0,n}(dy^n)\\
&\quad + (1-\lambda)\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}^2_{0,n})(dx^n,dy^n)\\
&\quad - (1-\lambda)\int_{\mathcal{Y}_{0,n}} \log\Big(\frac{\nu^2_{0,n}(dy^n)}{U_{0,n}(dy^n)}\Big)\nu^2_{0,n}(dy^n).
\end{aligned}$$

Finally, since $\nu^1_{0,n}(\cdot)\ll U_{0,n}(\cdot)$ and $\nu^2_{0,n}(\cdot)\ll U_{0,n}(\cdot)$ by substituting the following versions $\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)\times\nu^i_{0,n}(dy^n)}{U_{0,n}(dy^n)\times\nu^i_{0,n}(dy^n)}$, $i=1,2$, of the RND for $\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{U_{0,n}(dy^n)}$ in the first and third right hand side expression in the preceding equations yields

$$\mathbb{I}_{X^n\to Y^n}\big(\lambda\overleftarrow{P}^1_{0,n}+(1-\lambda)\overleftarrow{P}^2_{0,n},\overrightarrow{Q}_{0,n}\big)\geq\lambda\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}^1_{0,n},\overrightarrow{Q}_{0,n}\big)+(1-\lambda)\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}^2_{0,n},\overrightarrow{Q}_{0,n}\big).$$

This completes the derivation of 2).

3) Here, it will be shown that for $\overrightarrow{Q}^1_{0,n}(\cdot|x^n)$, $\overrightarrow{Q}^2_{0,n}(\cdot|x^n)\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ such that $\overrightarrow{Q}^1_{0,n}(\cdot|x^n)\neq\overrightarrow{Q}^2_{0,n}(\cdot|x^n)$, and $\lambda\in(0,1)$, then $\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\lambda\overrightarrow{Q}^1_{0,n}+(1-\lambda)\overrightarrow{Q}_{0,n}\big)<\lambda\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\overrightarrow{Q}^1_{0,n}\big)+(1-\lambda)\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\overrightarrow{Q}^2_{0,n}\big)$, for a fixed $\overleftarrow{P}_{0,n}\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$.

It is already known that $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})$ is a convex functional on $\overrightarrow{Q}_{0,n}\in\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$ for a fixed $\overleftarrow{P}_{0,n}\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$. All is required to show in order to have strict convexity is that $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})<\infty$. This can be easily obtained from part 1) since $\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n}\ll\overleftarrow{P}_{0,n}\otimes\nu_{0,n}$ if and only if $\overrightarrow{Q}_{0,n}(\cdot|x^n)\ll\nu_{0,n}(\cdot)$, for $\mu_{0,n}-$almost all $x^n\in\mathcal{X}_{0,n}$. Hence, from the strict convexity of the function $s\log s$, $s\in[0,\infty)$, and the expression of directed information as a functional of $\{\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n}\}\in\mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\times\mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n};\mathcal{X}_{0,n})$, with $\overrightarrow{Q}_{0,n}=\lambda\overrightarrow{Q}^1_{0,n}+(1-\lambda)\overrightarrow{Q}^2_{0,n}$ it follows that

$$\begin{aligned}
\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n},\overrightarrow{Q}_{0,n})&=\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\Big(\frac{(\overleftarrow{P}_{0,n}\otimes\overrightarrow{Q}_{0,n})(dx^n,dy^n)}{\overrightarrow{\Pi}(dx^n,dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&=\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overrightarrow{Q}_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&\leq\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\lambda\log\Big(\frac{\overrightarrow{Q}^1_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overrightarrow{Q}^1_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&+\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}(1-\lambda)\log\Big(\frac{\overrightarrow{Q}^2_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overrightarrow{Q}^2_{0,n}\otimes\overleftarrow{P}_{0,n})(dx^n,dy^n)\\
&=\lambda\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\overrightarrow{Q}^1_{0,n}\big)+(1-\lambda)\mathbb{I}_{X^n\to Y^n}\big(\overleftarrow{P}_{0,n},\overrightarrow{Q}^2_{0,n}\big)\\
&<\infty.
\end{aligned}$$

This completes the derivation of 3).

C. Proof of Theorem III.5

A1) First, it is shown that the joint distribution of the basic joint process $\{(X^{(\alpha)}_i,Y^{(\alpha)}_i):i\in\mathbb{N}\}$ converges as $\alpha\longrightarrow\infty$ to the joint distribution of a joint process $\{(X^{(0)}_i,Y^{(0)}_i):$

$i \in \mathbb{N}\}$ and secondly, that this limiting joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ is also a basic joint process corresponding to the backward channel $\overleftarrow{P}_{0,n}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$, that is, $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{\alpha}_{0,n})(dx^n, dy^n) \xrightarrow{w} (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{0}_{0,n})(dx^n, dy^n) \in \mathcal{M}_1(\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n})$ and that $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}^{0}_{0,n})(dx^n, dy^n)$ has backward channel $\overleftarrow{P}_{0,n}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$.

The first part is established by showing weak compactness of the family of measures induced by $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$. According to Prohorov's theorem (Appendix VI-A, Theorem VI.4), the family of joint probability distributions of the sequence $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : \ i \in \mathbb{N}\}$ is weakly compact if it is tight (uniformly tight). Therefore, we shall show tightness of the family of measures induced by the joint sequence of RV's $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$.

Utilizing condition $\mathbf{CA}$ for a fixed $n \in \mathbb{N}$, for any compact sets $K_0 \in \mathcal{X}_0, K_1 \in \mathcal{X}_1, \ldots, K_{n-1} \in \mathcal{X}_{n-1}$ the family of distributions $\{p_n(\cdot; x^{n-1}, y^{n-1}) : x_0 \in K_0, x_1 \in K_1, \ldots, x_{n-1} \in K_{n-1}, y^{n-1} \in \mathcal{Y}_{0,n-1}\} \subset \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1})$ is compact. Indeed, given any joint sequence $\{X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}\}$ by selecting a subsequence α_i such that the subsequence $\{X_0^{(\alpha_i)}, \ldots, X_{n-1}^{(\alpha_i)}, Y_0^{(\alpha_i)}, \ldots, Y_{n-1}^{(\alpha_i)}\}$ converges to $\{X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)}\}$, a weakly convergent subsequence of measures $p_n(\cdot; x_0^{(\alpha_i)}, \ldots, x_{n-1}^{(\alpha_i)}, y_0^{(\alpha_i)}, \ldots, y_{n-1}^{(\alpha_i)})$ is obtained. By Prohorov's theorem, a family of measures on a complete separable metric spaces is weakly compact if and only if it is tight. Hence, for any sequence of compact sets $K_0 \subset \mathcal{X}_0, K_1 \subset \mathcal{X}_1, \ldots, K_{n-1} \subset \mathcal{X}_{n-1}$, and $\epsilon_1 > 0$ a compact set $K_n \subset \mathcal{X}_n$ can be constructed such that $p_n(K_n; x^{n-1}, y^{n-1}) \geq 1 - \epsilon_1$, for any $x_0 \in K_0, x_1 \in K_1, \ldots, x_{n-1} \in K_{n-1}, y^{n-1} \in \mathcal{Y}_{0,n-1}$. To this end, pick $\epsilon_1 > 0$ and construct the compact sets as follows. Choose compact set $K_0 \subset \mathcal{X}_0$ such that $p_0(K_0) \geq 1 - \frac{\epsilon_1}{2}$, compact set $K_1 \subset \mathcal{X}_1$ such that $p_1(K_1; x_0, y_0) \geq 1 - \frac{\epsilon_1}{2^2}$, for any $x_0 \in K_0, y_0 \in \mathcal{Y}_0$, compact set $K_2 \subset \mathcal{X}_2$ such that $p_2(K_2; x_0, x_1, y_0, y_1) \geq 1 - \frac{\epsilon_1}{2^3}$, for any $x_0 \in K_0, x_1 \in K_1, y_0 \in \mathcal{Y}_0, y_1 \in \mathcal{Y}_1$, and compact set K_n such that $p_n(K_n; x_0, \ldots, x_{n-1}, y_0, \ldots, y_{n-1}) \geq 1 - \frac{\epsilon_1}{2^{n+1}}$, for any $x_0 \in K_0, \ldots, x_{n-1} \in K_{n-1}, y^{n-1} \in \mathcal{Y}_{0,n-1}$.

Likewise, since $\{\mathcal{Y}_i : i \in \mathbb{N}\}$ are compact Polish spaces, and measures on compact Polish spaces are weakly compact (Appendix VI-A, Theorem VI.1), then the family of conditional distributions $\{q_n(\cdot; y^{n-1}, x^n) : y^{n-1} \in \mathcal{Y}_{0,n-1}, x^n \in \mathcal{X}_{0,n}\} \subset \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n})$ is weakly compact. Utilizing the condition of weak compactness of a family of measures we verify that for any sequence of compact sets $\Phi_0 \subset \mathcal{Y}_0, \Phi_1 \subset \mathcal{Y}_1, \ldots, \Phi_{n-1} \subset \mathcal{Y}_{n-1}$, and $\epsilon_2 > 0$ there exists a compact set $\Phi_n \subset \mathcal{Y}_n$ such that $q_n(\Phi_n; y^{n-1}, x^n) \geq 1 - \epsilon_2$, for any $y_0 \in \Phi_0, y_1 \in \Phi_1, \ldots, y_{n-1} \in \Phi_{n-1}, x^n \in \mathcal{X}_{0,n}$. Choose an $\epsilon_2 > 0$ and construct compacts $\Phi_n \subset \mathcal{Y}_n$ as before: $\Phi_0 \subset \mathcal{Y}_0$ is such that $q_0(\Phi_0; x_0) \geq$

$1-\frac{\epsilon_2}{2}$, for any $x_0 \in \mathcal{X}_0$, and after $\Phi_0 \subset \mathcal{Y}_0, \Phi_1 \subset \mathcal{Y}_1, \ldots, \Phi_{n-1} \subset \mathcal{Y}_{n-1}$ are selected we choose $\Phi_n \subset \mathcal{Y}_n$ so that $q_n(\Phi_n; y^{n-1}, x^n) \geq 1 - \frac{\epsilon_2}{2^{n+1}}$, for any $y_0 \in \Phi_0, \ldots, y_{n-1} \in \Phi_{n-1}, x^n \in \mathcal{X}_{0,n}$.

Based on the above construction, which states that for any $n \in \mathbb{N}$ the families $p_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{X}_n; \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1})$ and $q_n(\cdot;\cdot,\cdot) \in \mathcal{Q}(\mathcal{Y}_n; \mathcal{Y}_{0,n-1} \times \mathcal{X}_{0,n})$ are tight, specifically,

$$p_n(K_n; x^{n-1}, y^{n-1}) \geq 1 - \frac{\epsilon_1}{2^{n+1}}, \quad q_n(\Phi_n; y^{n-1}, x^n) \geq 1 - \frac{\epsilon_2}{2^{n+1}}, \tag{VII.3}$$

for any $x_0 \in K_0, \ldots, x_n \in K_n$ and $y_0 \in \Phi_0, \ldots, y_{n-1} \in \Phi_{n-1}$, we show uniform tightness of the family of joint probability distributions of the joint sequence $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$. Under the above choice of $\{(K_i, \Phi_i) \subset \mathcal{X}_i \times \mathcal{Y}_i : i = 0, 1, \ldots, n\}$, we have the following inequalities.

$$\begin{aligned}
&\mathbb{P}\Big\{X_0^{(\alpha)} \in K_0, \ldots, X_n^{(\alpha)} \in K_n, Y_0^{(\alpha)} \in \Phi_0, \ldots, Y_n^{(\alpha)} \in \Phi_n\Big\} \\
&= \int_{\times_{i=0}^{n-1}(K_i \times \Phi_i)} \Bigg(\int_{K_n} p_n(dx; X_0^{(\alpha)} = x_0, Y_0^{(\alpha)} = y_0, \ldots, X_{n-1}^{(\alpha)} = x_{n-1}, Y_{n-1}^{(\alpha)} = y_{n-1}) \\
&\int_{\Phi_n} q_n(dy; Y_0^{(\alpha)} = y_0, X_0^{(\alpha)} = x_0, \ldots, Y_{n-1}^{(\alpha)} = y_{n-1}, X_n^{(\alpha)} = x) \Bigg) \mathbb{P}\Big\{X_0^{(\alpha)} \in dx_0, \ldots, X_{n-1}^{(\alpha)} \in dx_{n-1}, \\
&Y_0^{(\alpha)} \in dy_0, \ldots, Y_{n-1}^{(\alpha)} \in dy_{n-1}\Big\} \\
&\overset{(a)}{\geq} \left(1 - \frac{\epsilon_1}{2^{n+1}}\right)\left(1 - \frac{\epsilon_2}{2^{n+1}}\right) \mathbb{P}\Big\{X_0^{(\alpha)} \in K_0, \ldots, X_{n-1}^{(\alpha)} \in K_{n-1}, Y_0^{(\alpha)} \in \Phi_0, \ldots, Y_{n-1}^{(\alpha)} \in \Phi_{n-1}\Big\} \\
&= \left(1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_2}{2^{n+1}} + \frac{\epsilon_1\epsilon_2}{2^{2n+2}}\right) \mathbb{P}\Big\{X_0^{(\alpha)} \in K_0, \ldots, X_{n-1}^{(\alpha)} \in K_{n-1}, Y_0^{(\alpha)} \in \Phi_0, \ldots, Y_{n-1}^{(\alpha)} \in \Phi_{n-1}\Big\} \\
&\geq \left(1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_2}{2^{n+1}}\right) \mathbb{P}\Big\{X_0^{(\alpha)} \in K_0, \ldots, X_{n-1}^{(\alpha)} \in K_{n-1}, Y_0^{(\alpha)} \in \Phi_0, \ldots, Y_{n-1}^{(\alpha)} \in \Phi_{n-1}\Big\}
\end{aligned}$$

The inequality (a) follows from (VII.3). By repeating the procedure (or by perfect induction) we obtain the following inequalities.

$$\begin{aligned}
&\mathbb{P}\Big\{X_0^{(\alpha)} \in K_0, \ldots, X_n^{(\alpha)} \in K_n, Y_0^{(\alpha)} \in \Phi_0, \ldots, Y_n^{(\alpha)} \in \Phi_n\Big\} \\
&\qquad \geq 1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_2}{2^{n+1}} - \frac{\epsilon_1}{2^n} - \frac{\epsilon_2}{2^n} - \frac{\epsilon_1}{2^{n-1}} - \frac{\epsilon_2}{2^{n-1}} - \ldots - \frac{\epsilon_1}{2^1} - \frac{\epsilon_2}{2^1} \\
&\qquad = 1 - (\epsilon_1 + \epsilon_2) \sum_{i=1}^{n} \frac{1}{2^{i+1}} \geq 1 - (\epsilon_1 + \epsilon_2) \geq 1 - \epsilon, \qquad \text{for } \alpha = 1, 2, \ldots, \ n \in \mathbb{N}.
\end{aligned}$$

Hence, we have shown that the family of measures corresponding to the basic joint process $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$ is tight (uniformly), hence it converges weakly to the measures corresponding to the joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$. Thus, $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$ converges

in distribution to $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$, e.g., the basic joint process converges in distribution. Next, we show that the limiting joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ is a basic joint process with the same backward channel $\overleftarrow{P}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$. For any $n \in \mathbb{N}$, consider bounded and continuous real-valued functions $g_n(\cdot) \in BC(\mathcal{X}_n)$ and $\Psi_{0,n-1}(\cdot,\cdot) \in BC(\mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1})$. By the weak convergence of the joint measures corresponding to $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$ to the joint measures corresponding to $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ denoted by $(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n}^{\alpha})(dx^n, dy^n) \xrightarrow{w} (\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n}^{0})(dx^n, dy^n)$, the continuity of $g_n(\cdot)$ and the continuity of the function mapping $(x^{n-1}, y^{n-1}) \in \mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1} \longmapsto \int_{\mathcal{X}_n} g_n(x) p_n(dx; x^{n-1}, y^{n-1}) \in \mathbb{R}$, given $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that for all $\alpha \geq N$

$$\left| \int_{\mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1}} \left(\int_{\mathcal{X}_n} g_n(x) p_n(dx; x^{n-1}, y^{n-1}) \right) \Psi_{0,n-1}(x^{n-1}, y^{n-1}) P_{0,n-1}^{0}(dx^{n-1}, dy^{n-1}) \right.$$
$$\left. - \int_{\mathcal{X}_{0,n-1} \times \mathcal{Y}_{0,n-1}} \left(\int_{\mathcal{X}_n} g_n(x) p_n(dx; x^{n-1}, y^{n-1}) \right) \Psi_{0,n-1}(x^{n-1}, y^{n-1}) P_{0,n-1}^{\alpha}(dx^{n-1}, dy^{n-1}) \right| \leq \epsilon.$$

Since $\epsilon > 0$ is arbitrary, then

$$\lim_{\alpha \to \infty} \mathbb{E}\Big\{ g_n(X_n^{(\alpha)}) \Psi(X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Big\}$$
$$= \mathbb{E}\Big\{ g_n(X_n^{(0)}) \Psi(X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)}) \Big\}. \tag{VII.4}$$

Moreover, for all $\alpha = 1, 2, \ldots$, then

$$\mathbb{E}\Big\{ g_n(X_n^{(\alpha)}) \Psi(X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Big\}$$
$$= \mathbb{E}\Big\{ \Psi(X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \mathbb{E}\Big\{ g_n(X_n^{(\alpha)}) | X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Big\} \Big\}$$
$$= \mathbb{E}\Big\{ \Big(\int_{\mathcal{X}_n} g_n(x) p_n(dx; X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Big) \Psi(X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Big\}.$$

Hence, (VII.4) is equivalent to

$$\lim_{\alpha \to \infty} \mathbb{E}\Big\{ \int_{\mathcal{X}_n} g_n(x) p_n(dx; X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Psi(X_0^{(\alpha)}, \ldots, X_{n-1}^{(\alpha)}, Y_0^{(\alpha)}, \ldots, Y_{n-1}^{(\alpha)}) \Big\}$$
$$= \mathbb{E}\Big\{ \int_{\mathcal{X}_n} g_n(x) p_n(dx; X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{0}) \Psi(X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)}) \Big\}.$$

From the previous equality, which also holds for $\alpha = 0$, it follows that the next formula holds almost surely.

$$\mathbb{E}\Big\{ g_n(X_n^{(0)}) | X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)} \Big\}$$
$$= \int_{\mathcal{X}_n} g_n(x) p_n(dx; X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)}) - a.s.$$

Letting $g_n(\cdot)$ be the indicator function I_E, $E \in \mathcal{B}(\mathcal{X}_n)$ then

$$\mathbb{P}\Big\{ X_n^{(0)} \in E | X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)} \Big\} = p_n(E | X_0^{(0)}, \ldots, X_{n-1}^{(0)}, Y_0^{(0)}, \ldots, Y_{n-1}^{(0)}).$$

This shows that the limiting joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ is a basic process corresponding to the backward channel $\overleftarrow{P}_{0,n}(\cdot|\cdot) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$. This completes the derivation of A1).

A2) Here, we show that the marginal distributions generated by the joint sequence of RV's $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$ are weakly compact. In A1) it is shown that for any $n \in \mathbb{N}$, given $\epsilon > 0$, there are compact sets $K_0 \subset \mathcal{X}_0, K_1 \subset \mathcal{X}_1, \ldots, K_n \subset \mathcal{X}_n$ such that for any $x_0 \in K_0, x_1 \in K_1, \ldots, x_{n-1} \in K_{n-1}, y^{n-1} \in \mathcal{Y}_{0,n-1}$,

$$p_n(K_n; x^{n-1}, y^{n-1}) \geq 1 - \frac{\epsilon_1}{2^{n+1}}. \tag{VII.5}$$

Utilizing (VII.5) then

$$\mathbb{P}\Big\{ X_0^{(\alpha)} \in K_0, \ldots, X_n^{(\alpha)} \in K_n \Big\} = \mathbb{P}\Big\{ X_0^{(\alpha)} \in K_0, \ldots, X_n^{(\alpha)} \in K_n, Y_0^{(\alpha)} \in \mathcal{Y}_0, \ldots, Y_{n-1}^{(\alpha)} \in \mathcal{Y}_{n-1} \Big\}$$
$$= \int_{\times_{i=0}^n K_i} \int_{\mathcal{Y}_{0,n-1}} \mathbb{P}\Big\{ X_n^{(\alpha)} \in K_n | X_0^{(\alpha)} = x_0, \ldots, X_{n-1}^{(\alpha)} = x_{n-1}, Y_0^{(\alpha)} = y_0, \ldots, Y_{n-1}^{(\alpha)} = y_{n-1} \Big\}$$
$$\mathbb{P}\Big\{ X_0^{(\alpha)} \in dx_0, \ldots, X_{n-1}^{(\alpha)} \in dx_{n-1}, Y_0^{(\alpha)} \in dy_0, \ldots, Y_{n-1}^{(\alpha)} \in dy_{n-1} \Big\}$$
$$\geq \left(1 - \frac{\epsilon_1}{2^{n+1}}\right) \int_{\times_{i=0}^{n-1} K_i} \mathbb{P}\Big\{ X_0^{(\alpha)} \in dx_0, \ldots, X_{n-1}^{(\alpha)} \in dx_{n-1} \Big\}$$
$$= \left(1 - \frac{\epsilon_1}{2^{n+1}}\right) \mathbb{P}\Big\{ X_0^{(\alpha)} \in K_0, \ldots, X_{n-1}^{(\alpha)} \in K_{n-1} \Big\}$$
$$\geq \left(1 - \frac{\epsilon_1}{2^{n+1}}\right)\left(1 - \frac{\epsilon_1}{2^n}\right) \mathbb{P}\Big\{ X_0^{(\alpha)} \in K_0, \ldots, X_{n-2}^{(\alpha)} \in K_{n-2} \Big\}$$
$$= \left(1 - \frac{\epsilon_1}{2^{n+1}} - \frac{\epsilon_1}{2^n} + \frac{\epsilon_1^2}{2^{2n+1}}\right) \mathbb{P}\Big\{ X_0^{(\alpha)} \in K_0, \ldots, X_{n-2}^{(\alpha)} \in K_{n-2} \Big\}$$

$$\geq \left(1-\frac{\epsilon_1}{2^{n+1}}-\frac{\epsilon_1}{2^n}\right)\mathbb{P}\Big\{X_0^{(\alpha)}\in K_0,\ldots,X_{n-2}^{(\alpha)}\in K_{n-2}\Big\}$$
$$\geq \left(1-\frac{\epsilon_1}{2^{n+1}}-\frac{\epsilon_1}{2^n}\right)\left(1-\frac{\epsilon_1}{2^{n-1}}\right)\mathbb{P}\Big\{X_0^{(\alpha)}\in K_0,\ldots,X_{n-3}^{(\alpha)}\in K_{n-3}\Big\}$$
$$= \left(1-\frac{\epsilon_1}{2^{n+1}}-\frac{\epsilon_1}{2^n}-\frac{\epsilon_1}{2^{n-1}}+\frac{\epsilon_1^2}{2^{2n}}+\frac{\epsilon_1^2}{2^{2n-1}}\right)\mathbb{P}\Big\{X_0^{(\alpha)}\in K_0,\ldots,X_{n-3}^{(\alpha)}\in K_{n-3}\Big\}$$
$$\geq \left(1-\frac{\epsilon_1}{2^{n+1}}-\frac{\epsilon_1}{2^n}-\frac{\epsilon_1}{2^{n-1}}\right)\mathbb{P}\Big\{X_0^{(\alpha)}\in K_0,\ldots,X_{n-3}^{(\alpha)}\in K_{n-3}\Big\}.$$

By repeating the above procedure we obtain

$$\begin{aligned}\mathbb{P}\Big\{X_0^{(\alpha)}\in K_0,\ldots,X_n^{(\alpha)}\in K_n\Big\} &\geq 1-\frac{\epsilon_1}{2^{n+1}}-\frac{\epsilon_1}{2^n}-\frac{\epsilon_1}{2^{n-1}}-\ldots-\frac{\epsilon_1}{2^1}\\ &= 1-\epsilon_1\sum_{i=1}^{n}\frac{1}{2^{i+1}}\\ &\geq 1-\epsilon_1, \qquad \text{for all } \alpha=1,2,\ldots,n\in\mathbb{N}.\end{aligned}$$

This shows that the family of marginal distributions of the joint process $\{(X_i^{(\alpha)},Y_i^{(\alpha)}):i\in\mathbb{N}\}$ on $\mathcal{X}_{0,n}$ is uniformly tight and hence weakly compact. Consequently, the family of marginal distributions of the joint basic process $\{(X_i^{(\alpha)},Y_i^{(\alpha)}):i\in\mathbb{N}\}$ on $\mathcal{X}_{0,n}$ converges weakly to the family of joint process $\{(X_i^{(0)},Y_i^{(0)}):i\in\mathbb{N}\}$ on $\mathcal{X}_{0,n}$. Following the same methodology as before, it can be shown that the family of marginal distributions of the joint sequence $\{(X_i^{(\alpha)},Y_i^{(\alpha)}):i\in\mathbb{N}\}$ on $\mathcal{Y}_{0,n}$ is uniformly tight and hence weakly compact. Indeed, given $\epsilon_2>0$ there are compact sets $\Phi_0\subset\mathcal{Y}_0,\Phi_1\subset\mathcal{Y}_1,\ldots,\Phi_n\subset\mathcal{Y}_n$ such that

$$\begin{aligned}\mathbb{P}\Big\{y_0^{(\alpha)}\in\Phi_0,\ldots,y_n^{(\alpha)}\in\Phi_n\Big\} &\geq \left(1-\frac{\epsilon_2}{2^{n+1}}\right)\mathbb{P}\Big\{y_0^{(\alpha)}\in\Phi_0,\ldots,y_{n-1}^{(\alpha)}\in\Phi_{n-1}\Big\}\\ &\geq 1-\frac{\epsilon_2}{2^{n+1}}-\frac{\epsilon_2}{2^n}-\frac{\epsilon_2}{2^{n-1}}-\ldots-\frac{\epsilon_2}{2^1}\\ &= 1-\epsilon_2\sum_{i=1}^{n}\frac{1}{2^{i+1}}\\ &\geq 1-\epsilon_2, \text{ for all } \alpha=1,2,\ldots,n\in\mathbb{N}.\end{aligned}$$

This shows that the family of marginal distributions of the joint process $\{(X_i^{(\alpha)},Y_i^{(\alpha)}):i\in\mathbb{N}\}$ on $\mathcal{Y}_{0,n}$ is uniformly tight and hence weakly compact. Consequently, the family of marginal distributions of the joint process $\{(X_i^{(\alpha)},Y_i^{(\alpha)}):i\in\mathbb{N}\}$ on $\mathcal{Y}_{0,n}$ converges weakly to the family

of joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ on $\mathcal{Y}_{0,n}$. It was shown earlier that the basic process $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$ converges weakly to the joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ and that this limiting joint process has a same backward channel $\overleftarrow{P}_{0,n}(\cdot|y^{n-1})$ and a forward channel $\overrightarrow{Q}^0_{0,n}(\cdot|x^n)$. Hence, the marginal distributions of the basic joint process $\{(X_i^{(\alpha)}, Y_i^{(\alpha)}) : i \in \mathbb{N}\}$ converge to the marginal distributions of the basic joint process $\{(X_i^{(0)}, Y_i^{(0)}) : i \in \mathbb{N}\}$ corresponding to the backward channel $\overleftarrow{P}_{0,n}(\cdot|y^{n-1}) \in \mathcal{Q}^{\mathbf{C1}}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$ and a forward channel $\overrightarrow{Q}^0_{0,n}(\cdot|x^n) \in \mathcal{Q}^{\mathbf{C2}}(\mathcal{Y}_{0,n}; \mathcal{X}_{0,n})$. This completes the derivation of A2).

D. Proof of Theorem III.9

To show continuity of $\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$ with respect to weak convergence, we need to show that for every sequence $\{\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1}) : \alpha = 1, 2, \ldots\}$ such that $\overleftarrow{P}^{\alpha}_{0,n} \xrightarrow{w} \overleftarrow{P}^{0}_{0,n}$, we have

$$\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}^{\alpha}_{0,n}, \overrightarrow{Q}_{0,n}) \longrightarrow \mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}^{0}_{0,n}, \overrightarrow{Q}_{0,n}).$$

The derivation is based on the procedure utilized in [36] to show continuity for single letter mutual information. First, decompose the directed information into two terms as follows.

$$\begin{aligned}
\mathbb{I}_{X^n \to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n}) &= \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\nu_{0,n}(dy^n)}\Big)(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) \\
&= \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \log\Big(\frac{\overrightarrow{Q}_{0,n}(dy^n|x^n)}{\bar{\nu}_{0,n}(dy^n)}\Big)(\overleftarrow{P}_{0,n} \otimes \overrightarrow{Q}_{0,n})(dx^n, dy^n) \\
&- \int_{\mathcal{Y}_{0,n}} \log\Big(\frac{\nu_{0,n}(dy^n)}{\bar{\nu}_{0,n}(dy^n)}\Big)\nu_{0,n}(dy^n) \\
&= \int_{\mathcal{X}_{0,n} \times \mathcal{Y}_{0,n}} \Big(\xi_{\bar{\nu}_{0,n}}(x^n, y^n) \log \xi_{\bar{\nu}_{0,n}}(x^n, y^n)\Big)\overleftarrow{P}_{0,n}(dx^n|y^{n-1}) \otimes \bar{\nu}_{0,n}(dy^n) \\
&- \int_{\mathcal{Y}_{0,n}} \Big(\xi_{\bar{\nu}_{0,n}, \overleftarrow{P}_{0,n}}(y^n) \log \xi_{\bar{\nu}_{0,n}, \overleftarrow{P}_{0,n}}(y^n)\Big)\bar{\nu}_{0,n}(dy^n), \qquad \text{(VII.6)}
\end{aligned}$$

where $\xi_{\bar{\nu}_{0,n}, \overleftarrow{P}_{0,n}}(y^n) \stackrel{\triangle}{=} \frac{\nu_{0,n}(\cdot)}{\bar{\nu}_{0,n}(\cdot)}(y^n)$ emphasizes the fact that this RND depends on $\overleftarrow{P}_{0,n}$. For now, assume that both terms in on the right hand side of the above formula are finite; the validity of this assumption will be established at the end. Thus, we only need to show that both terms are bounded and continuous in the weak sense over $\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n}; \mathcal{Y}_{0,n-1})$.

Continuity of the first term. Since $\overleftarrow{P}^{\alpha}_{0,n} \xrightarrow{w} \overleftarrow{P}^{0}_{0,n}$, by [40, Theorem A.5.8, p. 320], utilizing Lebesgue's dominated convergence theorem, we have $\bar{\nu}_{0,n} \otimes \overleftarrow{P}^{\alpha}_{0,n} \xrightarrow{w} \bar{\nu}_{0,n} \otimes \overleftarrow{P}^{0}_{0,n}$. Since $\xi_{\bar{\nu}_{0,n}}(x^n, y^n)$ is continuous, then so is $\xi_{\bar{\nu}_{0,n}}(x^n, y^n) \log \xi_{\bar{\nu}_{0,n}}(x^n, y^n)$. By hypothesis, $\xi_{\bar{\nu}_{0,n}}(x^n, y^n) \log \xi_{\bar{\nu}_{0,n}}(x^n, y^n)$

is uniformly integrable over $\big\{\bar{\nu}_{0,n} \otimes \overleftarrow{P}_{0,n} : \overleftarrow{P}_{0,n} \in \overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})\big\}$. Therefore, using Theorem VI.12, Appendix VI-A, we conclude that

$$\lim_{\alpha\to\infty}\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\bar{\nu}_{0,n}(dy^n)\otimes\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1})$$
$$=\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}} \xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\bar{\nu}_{0,n}(dy^n)\otimes\overleftarrow{P}^{0}_{0,n}(dx^n|y^{n-1}). \quad \text{(VII.7)}$$

This proves the continuity of the first term. The finiteness of the first term is obtained from uniform integrability as follows. For a given $\epsilon>0$ and sufficiently large $c>0$

$$\sup_{\overleftarrow{P}_{0,n}\in\overleftarrow{Q}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})}\Big\{\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\big|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\big|I_{\{|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)|\geq c\}}$$
$$\times\bar{\nu}_{0,n}(dy^n)\otimes\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1})$$
$$+\int_{\mathcal{X}_{0,n}\times\mathcal{Y}_{0,n}}\big|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\big|I_{\{|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)|<c\}}\bar{\nu}_{0,n}(dy^n)\otimes\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1})\Big\}$$
$$\leq\sup_{\overleftarrow{P}_{0,n}\in\overleftarrow{Q}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})}\Big\{\int_{\{|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)|\geq c\}}\big|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\big|$$
$$\times\bar{\nu}_{0,n}(dy^n)\otimes\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1})\Big\}$$
$$+\sup_{\overleftarrow{P}_{0,n}\in\overleftarrow{Q}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})}\Big\{\int_{\{|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)|<c\}}\big|\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\big|$$
$$\times\bar{\nu}_{0,n}(dy^n)\otimes\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1})\Big\}$$
$$\leq\epsilon+c.$$

Continuity of the second term. For a fixed $y^n\in\mathcal{Y}_{0,n}$, since $\xi_{\bar{\nu}_{0,n}}(x^n,y^n)$ is uniformly integrable over $\overleftarrow{\mathcal{Q}}^{cl}(\mathcal{X}_{0,n};\mathcal{Y}_{0,n-1})$, by Theorem VI.10, Appendix VI-A, we deduce that $\overleftarrow{P}^{\alpha}_{0,n}\xrightarrow{w}\overleftarrow{P}^{0}_{0,n}$ implies pointwise convergence of $\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\longrightarrow\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{0}_{0,n}}(y^n)$. By continuity of the logarithm, we obtain the pointwise convergence of $\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\log\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\longrightarrow\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{0}_{0,n}}(y^n)\log\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{0}_{0,n}}(y^n)$. It only remains to show convergence under the integral with respect to $\bar{\nu}_{0,n}$. By (III.34), then $\forall\alpha$

$$\Big|\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\log\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\Big|\leq\frac{2}{e\ln 2}+\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\log\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)$$
$$=\frac{2}{e\ln 2}\int_{\mathcal{X}_{0,n}}\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\log\xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1}) \quad \text{(VII.8)}$$
$$\leq\frac{2}{e\ln 2}+\int_{\mathcal{X}_{0,n}}\Big(\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\log\xi_{\bar{\nu}_{0,n}}(x^n,y^n)\Big)\overleftarrow{P}^{\alpha}_{0,n}(dx^n|y^{n-1}).$$

where (VII.8) follows from (VII.6) and the nonnegativity of $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$. By (VII.7), the integration of the right hand side over $\bar{\nu}_{0,n}$ converges. Thus, by the generalized Lebesgue's dominated convergence theorem [42, p. 59], we conclude that

$$\int_{\mathcal{Y}_{0,n}} \xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\log \xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{\alpha}_{0,n}}(y^n)\bar{\nu}_{0,n}(dy^n) \stackrel{\alpha\to\infty}{\longrightarrow} \int_{\mathcal{Y}_{0,n}} \xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{0}_{0,n}}(y^n)\log \xi_{\bar{\nu}_{0,n},\overleftarrow{P}^{0}_{0,n}}(y^n)\bar{\nu}_{0,n}(dy^n).$$

This implies the continuity of the second term. Furthermore, its finiteness follows as before. Since both terms are finite and continuous we deduce continuity of the directed information $\mathbb{I}_{X^n\to Y^n}(\overleftarrow{P}_{0,n}, \overrightarrow{Q}_{0,n})$. This completes the derivation.

ACKNOWLEDGEMENT

This work was financially supported by a medium size University of Cyprus grant entitled "DIMITRIS". Parts of the material in this paper were presented at the IEEE International Symposium on Information Theory, Boston MA, July 1–6 2012 [45], and submitted for presentation at the IEEE International Symposium on Information Theory, Istanbul, Turkey, July 7–12 2013 [46].

REFERENCES

[1] H. Marko, "The bidirectional communication theory–A generalization of information theory," *IEEE Transactions on Communications*, vol. 21, no. 12, pp. 1345–1351, Dec. 1973.

[2] J. L. Massey, "Causality, feedback and directed information," in *International Symposium on Information Theory and its Applications (ISITA '90)*, Nov. 27-30 1990, pp. 303–305.

[3] G. Kramer, "Directed information for channels with feedback," Ph.D. dissertation, Swiss Federal Institute of Technology (ETH), December 1998.

[4] S. C. Tatikonda, "Control over communication constraints," Ph.D. dissertation, Mass. Inst. of Tech. (M.I.T.), Cambridge, MA, 2000.

[5] J. Chen and T. Berger, "The capacity of finite-state Markov channels with feedback," *IEEE Transactions on Information Theory*, vol. 51, no. 3, pp. 780–798, Mar. 2005.

[6] S. Yang, A. Kavcic, and S. Tatikonda, "Feedback capacity of finite-state machine channels," *IEEE Transactions on Information Theory*, vol. 51, no. 3, pp. 799–810, Mar. 2005.

[7] H. Permuter, P. Cuff, B. Van Roy, and T. Weissman, "Capacity of the trapdoor channel with feedback," *IEEE Transactions on Information Theory*, vol. 54, no. 7, pp. 3150–3165, July 2008.

[8] S. Tatikonda and S. Mitter, "The capacity of channels with feedback," *IEEE Transactions on Information Theory*, vol. 55, no. 1, pp. 323–349, Jan. 2009.

[9] H. H. Permuter, T. Weissman, and J. Chen, "Capacity region of the finite-state multiple-access channel with and without feedback," *IEEE Transactions on Information Theory*, vol. 55, no. 6, pp. 2455–2477, June 2009.

[10] H. H. Permuter, T. Weissman, and A. J. Goldsmith, "Finite state channels with time-invariant deterministic feedback," *IEEE Transactions on Information Theory*, vol. 55, no. 2, pp. 644–662, Feb. 2009.

[11] B. Shrader and H. Permuter, "Feedback capacity of the compound channel," *IEEE Transactions on Information Theory*, vol. 55, no. 8, pp. 3629–3644, Aug. 2009.

[12] R. Venkataramanan, "Information-theoretic results on communication problems with feed-forward and feedback," Ph.D. dissertation, University of Michigan-Ann Arbor, December 2008.

[13] G. Kramer, "Capacity results for the discrete memoryless network," *IEEE Transactions on Information Theory*, vol. 49, no. 1, pp. 4–21, Jan. 2003.

[14] H. H. Permuter, Y.-H. Kim, and T. Weissman, "Interpretations of directed information in portfolio theory, data compression, and hypothesis testing," *IEEE Transactions on Information Theory*, vol. 57, no. 6, pp. 3248–3259, 2011.

[15] V. Solo, "On causality and mutual information," in *47th IEEE Conference on Decision and Control (CDC '08)*, Dec. 2008, pp. 4939–4944.

[16] P. O. Amblard and O. J. J. Michel, "On directed information theory and granger causality graphs," *Journal of Computational Neuroscience*, vol. 30, no. 1, pp. 7–16, Feb. 2011.

[17] C. J. Quinn, T. P. Coleman, N. Kiyavash, and N. G. Hatsopoulos, "Estimating the directed information to infer causal relationships in ensemble neural spike train recordings," *Journal of Computational Neuroscience*, vol. 30, no. 1, pp. 17–44, Feb. 2011.

[18] T. S. Han and S. Verdu, "Approximation theory of output statistics," *IEEE Transactions on Information Theory*, vol. 39, no. 3, pp. 752–772, May 1993.

[19] R. E. Blahut, *Principles and Practice of Information Theory*, ser. in Electrical and Computer Engineering. Reading, MA: Addison-Wesley Publishing Company, 1987.

[20] I. Csiszár, "Arbitrarily varying channels with general alphabets and states," *IEEE Transactions on Information Theory*, vol. 38, no. 6, pp. 1725–1742, Nov. 1992.

[21] I. I. Gihman and A. V. Skorohod, *Controlled Stochastic Processes*. Springer-Verlag, New York Inc., 1979, translated by Samuel Kotz.

[22] D. P. Bertsekas and S. E. Shreve, *Stochastic Optimal Control: The Discrete-Time Case*, ser. Optimization and Neural Computation Series. Athena Scientific, 2007.

[23] S. Tatikonda and S. Mitter, "The capacity of channels with feedback," *IEEE Transactions on Information Theory*, vol. 55, no. 1, pp. 323–349, Jan. 2009.

[24] P. A. Stavrou, C. D. Charalambous, and C. K. Kourtellaris, "Optimal nonstationary reproduction distribution for nonanticipative rdf on abstract alphabets," *CoRR*, vol. abs/1301.6522, 2013. [Online]. Available: http://arxiv.org/abs/1301.6522

[25] G. Kramer, "Topics in multi-user information theory," *Foundations and Trends in Communications and Information Theory*, vol. 4, no. 4-5, pp. 265–444, Apr. 2007.

[26] E. A. Gamal and H. Y. Kim, *Network Information Theory*. Cambridge University Press, December 2011.

[27] I. Csiszár, "On an extremum problem of information theory," *Studia Scientiarum Mathematicarum Hungarica*, vol. 9, pp. 57–71, 1974.

[28] R. B. Ash and C. A. Doléans-Dade, *Probability and Measure Theory*, 2nd ed. Academic Press, December 1999.

[29] M. Pinsker, *Information and Information Stability of Random Variables and Processes*. Holden-Day Inc, San Francisco, 1964, translated by Amiel Feinstein.

[30] S. Ihara, *Information theory - for Continuous Systems*. World Scientific, 1993.

[31] A. N. Shiryaev, *Probability*, 2nd ed., ser. Graduate Texts in Mathematics. Springer-Verlag, Berlin, Heidelberg, New York, 1996.

[32] P. Billingsley, *Convergence of Probability Measures*, 2nd ed., ser. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., 1999.

[33] T. Berger, "Rate distortion theory for sources with abstract alphabets and memory," *Information and Control*, vol. 13, no. 3, pp. 254–273, Sep. 1968.

[34] A. Lapidoth and P. Narayan, "Reliable communication under channel uncertainty," *IEEE Transactions on Information Theory*, vol. 44, no. 6, pp. 2148–2177, Oct. 1998.

[35] D. J. Sakrison, "The rate distortion function for a class of sources," *Information and Control*, vol. 15, no. 2, pp. 165–195, Aug. 1969.

[36] M. Fozunbal, S. McLaughlin, and R. Schafer, "Capacity analysis for continuous-alphabet channels with side information, part I: A general framework," *IEEE Transactions on Information Theory*, vol. 51, no. 9, pp. 3075–3085, Sep. 2005.

[37] P. A. Stavrou and C. D. Charalambous, "Two extremum problems of directed information theory," *preprint*, 2013.

[38] F. Topsøe, "An information theoretical identity and a problem involving capacity," *Studia Scientiarum Mathematicarum Hungarica*, vol. 2, pp. 291–292, 1967.

[39] R. Blahut, "Computation of channel capacity and rate-distortion functions," *IEEE Transactions on Information Theory*, vol. 18, no. 4, pp. 460–473, July 1972.

[40] P. Dupuis and R. S. Ellis, *A Weak Convergence Approach to the Theory of Large Deviations*. John Wiley & Sons, Inc., New York, 1997.

[41] K. R. Parthasarathy, *Probability Measures on Metric Spaces*. Academic Press, New York, 1967.

[42] G. B. Folland, *Real Analysis: Modern Techniques and Their Applications*, 2nd ed. John Wiley & Sons Inc., April 1999.

[43] J. D. Deuschel and D. W. Stroock, *Large Deviations*. San Diego: Academic Press, Inc., 1989.

[44] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. John Wiley & Sons, Inc., Hoboken, New Jersey, 2006.

[45] C. D. Charalambous and P. A. Stavrou, "Directed information on abstract spaces: properties and extremum problems," in *IEEE International Symposium on Information Theory (ISIT)*, July 1-6 2012, pp. 518–522.

[46] P. A. Stavrou and C. D. Charalambous, "Variational equalities of directed information and applications," *CoRR*, vol. abs/1301.6520, 2013. [Online]. Available: http://arxiv.org/abs/1301.6520